




## INTERNATIONAL APPLICATION PUBLISHED UNDER THE PATENT COOPERATION TREATY (PCT)

(51) International Patent Classification <sup>6</sup> : <b>C07K 14/00</b>		A2	(11) International Publication Number: <b>WO 99/60021</b>
			(43) International Publication Date: 25 November 1999 (25.11.99)
(21) International Application Number: PCT/US99/10953		(81) Designated States: AE, AL, AM, AT, AU, AZ, BA, BB, BG, BR, BY, CA, CH, CN, CU, CZ, DE, DK, EE, ES, FI, GB, GD, GE, GH, GM, HR, HU, ID, IL, IN, IS, JP, KE, KG, KP, KR, KZ, LC, LK, LR, LS, LT, LU, LV, MD, MG, MK, MN, MW, MX, NO, NZ, PL, PT, RO, RU, SD, SE, SG, SI, SK, SL, TJ, TM, TR, TT, UA, UG, UZ, VN, YU, ZA, ZW, ARIPO patent (GH, GM, KE, LS, MW, SD, SL, SZ, UG, ZW), Eurasian patent (AM, AZ, BY, KG, KZ, MD, RU, TJ, TM), European patent (AT, BE, CH, CY, DE, DK, ES, FI, FR, GB, GR, IE, IT, LU, MC, NL, PT, SE), OAPI patent (BF, BJ, CF, CG, CI, CM, GA, GN, GW, ML, MR, NE, SN, TD, TG).	
(22) International Filing Date: 19 May 1999 (19.05.99)			
(30) Priority Data: 124550 19 May 1998 (19.05.98) IL PCT/US98/14715 21 July 1998 (21.07.98) US 09/218,277 22 December 1998 (22.12.98) US			
(71) Applicant: YEDA RESEARCH AND DEVELOPMENT CO. LTD. [IL/IL]; P.O. Box 95, 76100 Rehovot (IL).			
(71) Applicant (for SD only): MCINNIS, Patricia, A. [US/US]; Apartment #203, 2325 42nd Street N.W., Washington, DC 20007 (US).		Published Without international search report and to be republished upon receipt of that report.	
(72) Inventors: EISENBACH-SCHWARTZ, Michal; Rupin Street 5, 76353 Rehovot (IL). COHEN, Irun, R.; Hankin Street 11, 76343 Rehovot (IL). BESERMAN, Pierre; 76834 Moshav Sitriya (IL). MOSONEGO, Alon; Ben-Yosef, 73112 Kfar Hanoar Ben-Shemen (IL). MOALEM, Gila; Bosel Street 27, 76405 Rehovot (IL).			
(74) Agent: BROWDY, Roger, L.; Browdy and Neimark, P.L.L.C., Suite 300, 419 Seventh Street N.W., Washington, DC 20004 (US).			

(54) Title: ACTIVATED T CELLS, NERVOUS SYSTEM-SPECIFIC ANTIGENS AND THEIR USES

## (57) Abstract

Compositions and methods are provided for treating injury to or disease of the central or peripheral nervous system. In one embodiment, treatment is effected using activated T cells that recognize an antigen of the nervous system or a peptide derived therefrom or a derivative thereof to promote nerve regeneration or to prevent or inhibit neuronal degeneration within the nervous system. Treatment involves administering an NS-specific antigen or peptide derived therefrom or a derivative thereof, or a nucleotide sequence encoding said antigen or peptide, to promote nerve regeneration or to prevent or inhibit neuronal degeneration in the nervous system, either the central nervous system or the peripheral nervous system. The NS-specific activated T cells can be administered alone or in combination with NS-specific antigen or peptide derived therefrom or a derivative thereof or a nucleotide sequence encoding said antigen or peptide, or any combination thereof.



*FOR THE PURPOSES OF INFORMATION ONLY*

Codes used to identify States party to the PCT on the front pages of pamphlets publishing international applications under the PCT.

AL	Albania	ES	Spain	LS	Lesotho	SI	Slovenia
AM	Armenia	FI	Finland	LT	Lithuania	SK	Slovakia
AT	Austria	FR	France	LU	Luxembourg	SN	Senegal
AU	Australia	GA	Gabon	LV	Latvia	SZ	Swaziland
AZ	Azerbaijan	GB	United Kingdom	MC	Monaco	TD	Chad
BA	Bosnia and Herzegovina	GE	Georgia	MD	Republic of Moldova	TG	Togo
BB	Barbados	GH	Ghana	MG	Madagascar	TJ	Tajikistan
BE	Belgium	GN	Guinea	MK	The former Yugoslav Republic of Macedonia	TM	Turkmenistan
BF	Burkina Faso	GR	Greece	ML	Mali	TR	Turkey
BG	Bulgaria	HU	Hungary	MN	Mongolia	TT	Trinidad and Tobago
BJ	Benin	IE	Ireland	MR	Mauritania	UA	Ukraine
BR	Brazil	IL	Israel	MW	Malawi	UG	Uganda
BY	Belarus	IS	Iceland	MX	Mexico	US	United States of America
CA	Canada	IT	Italy	NZ	Niger	UZ	Uzbekistan
CF	Central African Republic	JP	Japan	NL	Netherlands	VN	Viet Nam
CG	Congo	KE	Kenya	NO	Norway	YU	Yugoslavia
CH	Switzerland	KG	Kyrgyzstan	NZ	New Zealand	ZW	Zimbabwe
CI	Côte d'Ivoire	KP	Democratic People's Republic of Korea	PL	Poland		
CM	Cameroon	KR	Republic of Korea	PT	Portugal		
CN	China	KZ	Kazakhstan	RO	Romania		
CU	Cuba	LC	Saint Lucia	RU	Russian Federation		
CZ	Czech Republic	LI	Liechtenstein	SD	Sudan		
DE	Germany	LK	Sri Lanka	SE	Sweden		
DK	Denmark	LR	Liberia	SG	Singapore		
EE	Estonia						

## ACTIVATED T CELLS, NERVOUS SYSTEM-SPECIFIC ANTIGENS AND THEIR USES

### Field of the Invention

The present invention relates to compositions and methods for the promotion of nerve regeneration or prevention or inhibition of neuronal degeneration to ameliorate the effects of injury or disease of the nervous system (NS). In certain embodiments, activated antiself T cells, an NS-specific antigen or peptide derived therefrom or a nucleotide sequence encoding an NS-specific antigen or peptide derived therefrom can be used to promote nerve regeneration or to prevent or inhibit neuronal degeneration caused by injury or disease of nerves within the central nervous system or peripheral nervous system of a human subject. The compositions of the present invention may be administered alone or may be optionally administered in any desired combination.

### Background of the Invention

The nervous system comprises the central (CNS) and the peripheral (PNS) nervous system. The central nervous system is composed of the brain and spinal cord; the peripheral nervous system consists of all of the other neural elements, namely the nerves and ganglia outside of the brain and spinal cord.

Damage to the nervous system may result from a traumatic injury, such as penetrating trauma or blunt trauma, or a disease or disorder, including but not limited to Alzheimer's disease, Parkinson's disease, multiple sclerosis, Huntington's disease, amyotrophic lateral sclerosis (ALS), diabetic neuropathy, senile dementia, and ischemia.

Maintenance of central nervous system integrity is a complex "balancing act" in which compromises are struck with the immune system. In most tissues, the immune system plays an essential part in protection, repair, and healing. In the central nervous system, because of its unique immune privilege, immunological reactions are relatively limited (Streilein, J.W., 1993, Curr. Opin. Immunol. 5:428-423; Streilein, J.W., Science 270:1158-1159). A growing body of evidence indicates

that the failure of the mammalian central nervous system to achieve functional recovery after injury is a reflection of an ineffective dialog between the damaged tissue and the immune system. For example, the restricted communication between the central nervous system and blood-borne macrophages affects the capacity of axotomized axons to regrow; transplants of activated macrophages can promote central nervous system regrowth (Lazarov Spiegler, O., et al., 1996, FASEB J. 10:1296-1302; Rapalino, O. et al., 1998, Nature Med. 4:814-821).

Activated T cells have been shown to enter the central nervous system parenchyma, irrespective of their antigen specificity, but only T cells capable of reacting with a central nervous system antigen seem to persist there (Hickey, W.F. et al., 1991, J. Neurosci. Res. 28:254-260; Werkele, H., 1993, In The Blood-Brain Barrier, Pardridge, Ed., Raven Press, Ltd. New York, 67-85; Kramer, R. et al., 1995, Nature Med. 1(11):1162-1166)). T cells reactive to antigens of central nervous system white matter, such as myelin basic protein (MBP), can induce the paralytic disease experimental autoimmune encephalomyelitis (EAE) within several days of their inoculation into naive recipient rats (Ben Nun, A., et al., 1981, Eur. J. Immunol. 11:195-199). Anti-MBP T cells may also be involved in the human disease multiple sclerosis (Ota, K. et al., 1990 Nature 346:183-187; Martin, R. 1997, J. Neural Transm. Suppl. 49:53-67). However, despite their pathogenic potential, anti-MBP T cell clones are present in the immune systems of healthy subjects (Burns, J., et al. 1983, Cell Immunol. 81:435-440; Pette, M. et al., 1990, Proc. Natl. Acad. Sci. USA 87:7968-7972; Martin, R. et al., 1990, J. Immunol. 145:540-548; Schiuesener, H.J, et al., 1985, J. Immunol. 135:3128-3133). Activated T cells, which normally patrol the intact central nervous system, transiently accumulate at sites of central nervous system white matter lesions (Hirschberg, D.L., et al., 1998, J. Neuroimmunol. 89:88-96).

A catastrophic consequence of central nervous system injury is that the primary damage is often compounded by the gradual secondary loss of adjacent neurons that apparently were undamaged, or only marginally damaged, by the initial injury



(Faden, A. I., et al., 1992, Trends Pharmacol. Sci. 13:29-35; Faden, A.I., 1993, Crit. Rev. Neurobiol. 7:175-186; McIntosh, T.K., 1993, J. Neurotrauma 10:215-261). The primary lesion causes changes in extracellular ion concentrations, elevation of amounts of free radicals, release of neurotransmitters, depletion of growth factors, and local inflammation. These changes trigger a cascade of destructive events in the adjacent neurons that initially escaped the primary injury (Lynch, D.R. et al., 1994, Curr. Opin. Neurol. 7:510-516; Bazan, N.G. et al., 1995, J. Neurotrauma 12:791-814; Wu, D. et al., 1994, J. Neurochem. 62:37-44). This secondary damage is mediated by activation of voltage-dependent or agonist-gated channels, ion leaks, activation of calcium-dependent enzymes such as proteases, lipases and nucleases, mitochondrial dysfunction and energy depletion, culminating in neuronal cell death (Yoshina, A. et al., 1991 Brain Res. 561:106-119; Hovda, D.A. et al., 1991, Brain Res. 567:1-10; Zivin, J.A., et al, 1991 Sci. Am. 265:56-63; Yoles, E. et al., 1992, Invest. Ophthalmol. Vis. Sci. 33:3586-3591). The widespread loss of neurons beyond the loss caused directly by the primary injury has been called "secondary degeneration."

Another tragic consequence of central nervous system injury is that neurons in the mammalian central nervous system do not undergo spontaneous regeneration following an injury. Thus, a central nervous system injury causes permanent impairment of motor and sensory functions.

Spinal cord lesions, regardless of the severity of the injury, initially result in a complete functional paralysis known as spinal shock. Some spontaneous recovery from spinal shock may be observed, starting a few days after the injury and tapering off within three to four weeks. The less severe the insult, the better the functional outcome. The extent of recovery is a function of the amount of undamaged tissue minus the loss due to secondary degeneration. Recovery from injury would be improved by neuroprotective treatment that could reduce secondary degeneration.

Citation or identification of any reference in this section or any other part of this application shall not be

construed as an admission that such reference is available as prior art to the invention.

#### SUMMARY OF THE INVENTION

The present invention is directed to methods and compositions for the promotion of nerve regeneration or prevention or inhibition of neuronal degeneration to ameliorate the effects of injury to or disease of the nervous system (NS). The present invention is based in part on the applicants' unexpected discovery that activated T cells that recognize an antigen of the NS of the patient promote nerve regeneration or confer neuroprotection. As used herein, "neuroprotection" refers to the prevention or inhibition of degenerative effects of injury or disease in the NS. Until recently, it was thought that the immune system excluded immune cells from participating in nervous system repair. It was quite surprising to discover that NS-specific activated T cells can be used to promote nerve regeneration or to protect nervous system tissue from secondary degeneration which may follow damage caused by injury or disease of the CNS or PNS.

"Activated T cell" as used herein includes (i) T cells that have been activated by exposure to a cognate antigen or peptide derived therefrom or derivative thereof and (ii) progeny of such activated T cells. As used herein, a cognate antigen is an antigen that is specifically recognized by the T cell antigen receptor of a T cell that has been previously exposed to the antigen. Alternatively, the T cell which has been previously exposed to the antigen may be activated by a mitogen, such as phytohemagglutinin (PHA) or concanavalin A.

In one embodiment, the present invention provides pharmaceutical compositions comprising a therapeutically effective amount of NS-specific activated T cells and methods for using such compositions to promote nerve regeneration or to prevent or inhibit neuronal degeneration in the CNS or PNS, in an amount which is effective to ameliorate the effects of an injury or disease of the NS. "NS-specific activated T cell" as used herein refers to an activated T cell having specificity for an antigen of the NS of a patient. The antigen used to confer the specificity to the T cells may be a self NS-antigen

of the patient, a peptide derived therefrom, or an NS-antigen of another individual or even another species, or a peptide derived therefrom, as long as the activated T cell recognizes an antigen in the NS of the patient.

The NS-specific activated T cells are used to promote nerve regeneration or to prevent or inhibit the effects of disease. If the disease being treated is an autoimmune disease, in which the autoimmune antigen is an NS antigen, the T cells which are used in accordance with the present invention for the treatment of neural damage or degeneration caused by such disease are preferably not activated against the same autoimmune antigen involved in the disease. While the prior art has described methods of treating autoimmune diseases by administering activated T cells to create a tolerance to the autoimmune antigen, the T cells of the present invention are not administered in such a way as to create tolerance, but are administered in such a way as to create accumulation of the T cells at the site of injury or disease so as to facilitate neural regeneration or to inhibit neural degeneration.

The prior art also discloses uses of immunotherapy against tumors, including brain tumors, by administering T cells specific to an NS antigen in the tumor so that such T cells may induce an immune system attack against the tumors. The present invention is not intended to comprehend such prior art techniques. However, the present invention is intended to comprehend the inhibition of neural degeneration or the enhancement of neural regeneration in patients with brain tumors by means other than the prior art immunotherapy of brain tumors. Thus, for example, NS-specific activated T cells, which are activated to an NS antigen of the patient other than an antigen which is involved in the tumor, would be expected to be useful for the purpose of the present invention and would not have been suggested by known immunotherapy techniques.

The present invention also provides pharmaceutical compositions comprising a therapeutically effective amount of an NS-specific antigen or peptide derived therefrom or derivative thereof and methods of use of such compositions to promote nerve regeneration or to prevent or inhibit neuronal degeneration in the CNS or PNS, in which the amount is

effective to activate T cells *in vivo* or *in vitro*, wherein the activated T cells inhibit or ameliorate the effects of an injury or disease of the NS. "NS-specific antigen" as used herein refers to an antigen that specifically activates T cells such that following activation the activated T cells accumulate at a site of injury or disease in the NS of the patient. In one embodiment, the peptide derived from an NS-specific antigen is a "cryptic epitope" of the antigen. A cryptic epitope activates specific T cells after an animal is immunized with the particular peptide, but not with the whole antigen. In another embodiment, the peptide derived from an NS-specific antigen is an immunogenic epitope of the antigen. "Derivatives" of NS-specific antigens or peptides derived therefrom as used herein refers to analogs or chemical derivatives of such antigens or peptides as described below, see Section 5.2.

The present invention also provides pharmaceutical compositions comprising a therapeutically effective amount of a nucleotide sequence encoding an NS-specific antigen or peptide derived therefrom or derivative thereof and methods of use of such compositions to promote nerve regeneration or for preventing or inhibiting neuronal degeneration in the CNS or PNS in which the amount is effective to ameliorate the effects of an injury or disease of the NS.

In the practice of the invention, therapy for amelioration of effects of injury or disease comprising administration of NS-specific activated T cells may optionally be in combination with an NS-specific antigen or peptide derived therefrom.

Additionally, oral administration of NS-specific antigen or a peptide derived therefrom, can be combined with active immunization to build up a critical T cell response immediately after injury.

In another embodiment cell banks can be established to store NS sensitized T cells for neuroprotective treatment of individuals at a later time, as needed. In this case, autologous T cells may be obtained from an individual. Alternatively, allogeneic or semi-allogeneic T cells may be stored such that a bank of T cells of each of the most common

MHC class II types are present. In case an individual is to be treated for an injury, preferably autologous stored T cells are used, but, if autologous T cells are not available, then cells should be used which share an MHC type II molecule with the patient, and these would be expected to be operable in that individual. The cells are preferably stored in an activated state after exposure to an NS antigen or peptide derived therefrom. However, the cells may also be stored in a resting state and activated once they are thawed and prepared for use. The cell lines of the bank are preferably cryopreserved. The cell lines are prepared in any way which is well known in the art. Once the cells are thawed, they are preferably cultured prior to injection in order to eliminate non-viable cells. During this culturing, the cells can be activated or reactivated using the same NS antigen or peptide as used in the original activation. Alternatively, activation may be achieved by culturing in the presence of a mitogen, such as phytohemagglutinin (PHA) or concanavalin A (preferably the former). This will place the cells into an even higher state of activation. The few days that it takes to culture the cells should not be detrimental to the patient as the treatment in accordance with the present invention may occur any time up to a week or more after the injury in order to still be effective. Alternatively, if time is of the essence, the stored cells may be administered immediately after thawing.

#### BRIEF DESCRIPTION OF THE FIGURES

Fig. 1 is a bar graph showing the presence of T cells in uninjured optic nerve or in injured optic nerve one week after injury. Adult Lewis rats were injected with activated T cells of the anti-MBP ( $T_{MBP}$ ), anti-OVA ( $T_{OVA}$ ), anti-p277 ( $T_{p277}$ ) lines, or with PBS, immediately after unilateral crush injury of the optic nerve. Seven days later, both the injured and uninjured optic nerves were removed, cryosectioned and analyzed immunohistochemically for the presence of immunolabeled T cells. T cells were counted at the site of injury and at randomly selected areas in the uninjured optic nerves. The histogram shows the mean number of T cells per  $mm^2 \pm$  s.e.m., counted in two to three sections of each nerve. Each

group contained three to four rats. The number of T cells was considerably higher in injured nerves of rats injected with anti-MBP, anti-OVA or anti-p277 T cells; statistical analysis (one-way ANOVA) showed significant differences between T cell numbers in injured optic nerves of rats injected with anti-MBP, anti-OVA, or anti-p277 T cells and the T cell numbers in injured optic nerves of rats injected with PBS ( $P < 0.001$ ); and between injured optic nerves and uninjured optic nerves of rats injected with anti-MBP, anti-OVA, or anti-p277 T cells ( $P < 0.001$ ).

Fig. 2 is a bar graph illustrating that T cells specific to MBP, but not of OVA or p277 or hsp60, protect neurons from secondary degeneration. Immediately after optic nerve injury, rats were injected with anti-MBP, anti-OVA or anti-p277 T cells, or with PBS. The neurotracer dye 4-Di-10-Asp was applied to optic nerves distal to the site of the injury, immediately after injury (for assessment of primary damage) or two weeks later (for assessment of secondary degeneration). Five days after dye application, the retinas were excised and flat-mounted. Labeled retinal ganglion cells (RGCs) from three to five randomly selected fields in each retina (all located at approximately the same distance from the optic disk) were counted by fluorescence microscopy. RGC survival in each group of injured nerves was expressed as the percentage of the total number of neurons spared after the primary injury (42% of neurons remained undamaged after the primary injury). The neuroprotective effect of anti-MBP T cells compared with that of PBS was significant ( $P < 0.001$ , one-way ANOVA). Anti-OVA T cells or anti-p277 T cells did not differ significantly from PBS in their effects on the protection of neurons that had escaped primary injury ( $P > 0.05$ , one-way ANOVA). The results are a summary of five experiments. Each group contained five to ten rats.

Figs. 3 (A-C) present photomicrographs of retrogradely labeled retinas of injured optic nerves of rats. Immediately after unilateral crush injury of their optic nerves, rats were injected with PBS (Fig. 3A) or with activated anti-p277 T cells (Fig. 3B) or activated anti-MBP T cells (Fig. 3C). Two weeks later, the neurotracer dye 4-Di-10-Asp was

applied to the optic nerves, distal to the site of injury. After 5 days, the retinas were excised and flat-mounted. Labeled (surviving) RGCs, located at approximately the same distance from the optic disk in each retina, were photographed.

Figs. 4(A-B) are graphs showing that clinical severity of EAE is not influenced by an optic nerve crush injury. For the results presented in Fig. 4A, Lewis rats, either uninjured (dash line) or immediately after optic nerve crush injury (solid line), were injected with activated anti-MBP T cells. EAE was evaluated according to a neurological paralysis scale. [Data points represent  $\pm$  s.e.m.] These results represent a summary of three experiments. Each group contained five to nine rats. Fig. 4B shows that the number of RGCs in the uninjured optic nerve is not influenced by injection of anti-MBP T cells. Two weeks after the injection of anti-MBP T cells or PBS, 4-Di-10Asp was applied to the optic nerves. After 5 days the retinas were excised and flat-mounted. Labeled RGCs from five fields (located at approximately the same distance from the optic disk) in each retina were counted and the average number per  $\text{mm}^2$  was calculated. There was no difference between the numbers of labeled RGCs in rats injected with anti-MBP T cells ( $T_{\text{MBP}}$ ) and in PBS-injected control rats.

Fig. 5 is a bar graph showing that T cells specific to p51-70 of MBP protect neurons from secondary degeneration. Immediately after optic nerve injury, rats were injected with anti-MBP T cells, anti-p51-70 T cells, or PBS. The neurotracer dye 4-Di-10-Asp was applied to optic nerves distal to the site of the injury, immediately after injury (for assessment of primary damage) or two weeks later (for assessment of secondary degeneration). Five days after dye application, the retinas were excised and flat-mounted. Labeled retinal ganglion cells (RGCs) from three to five randomly selected fields in each retina (all located at approximately the same distance from the optic disk) were counted by fluorescence microscopy. RGC survival in each group of injured nerves was expressed as the percentage of the total number of neurons spared after primary injury. Compared with

that of PBS treatment, the neuroprotective effects of anti-MBP anti-p51-70 T cells were significant ( $P < 0.001$ , one-way ANOVA).

Figs. 6(A-B) are graphs showing that anti-MBP T cells increase the compound action potential (CAP) amplitudes of injured optic nerves. Immediately after optic nerve injury, rats were injected with either PBS or activated anti-MBP T cells ( $T_{MBP}$ ). Two weeks later, the CAPs of injured (Fig. 6A) and uninjured (Fig. 6B) nerves were recorded. There were no significant differences in mean CAP amplitudes between uninjured nerves obtained from PBS-injected and T cell-injected rats ( $n=8$ ;  $p=0.8$ , Student's t-test). The neuroprotective effect of anti-MBP T cells (relative to PBS) on the injured nerve on day 14 after injury was significant ( $n=8$ ,  $p=0.009$ , Student's t-test).

Figs. 7(A-B) are graphs showing recovery of voluntary motor activity as a function of time after contusion, with and without injection of autoimmune anti-MBP T cells. (7A) Twelve rats were deeply anesthetized and laminectomized, and then subjected to a contusion insult produced by a 10 gram weight dropped from a height of 50 mm. Six of the rats, selected at random, were then inoculated i.p. with  $10^7$  anti-MBP T cells and the other six were inoculated with PBS. At the indicated time points, locomotor behavior in an open field was scored by observers blinded to the treatment received by the rats. Results are expressed as the mean values for each group. The vertical bars indicate SEM. Differences tested by repeated ANOVA, including all time points, were significant ( $p < 0.05$ ). (7B) A similar experiment using five PBS-treated animals and six animals treated with anti-MBP T cells were all subjected to a more severe contusion. At the indicated time points, locomotor behavior in an open field was scored. The results are expressed as the mean values for each group. The vertical bars indicate S.E.M. Rats in the treated group are represented by open circles and rats in the control group are represented by black circles. Horizontal bars show the median values. The inset shows the median plateau values of the two groups.

Figs 8(A-C) show retrograde labeling of cell bodies at the red nucleus in rats treated with autoimmune anti-MBP T cells (8A) and in control injured (8B) rats. Three months



after contusion and treatment with anti-MBP T cells, some rats from both the treated and the control groups were re-anesthetized and a dye was applied below the site of the contusion. After five to seven days the rats were again deeply anesthetized and their brains were excised, processed, and cryosectioned. Sections taken through the red nucleus were inspected and analyzed qualitatively and quantitatively under fluorescent and confocal microscopes. Significantly, more labelled nuclei were seen in the red nuclei of rats treated with anti-MBP T cells (8A) than in the red nuclei of PBS-treated rats (8B). The quantitative differences are shown in the bar graph (8C) and were obtained from animals with scores of 10 and 11 in the T cell treated group and scores of 6 in the control group. The bar graph shows mean  $\pm$  SD.

Fig. 9 is a series of photographs showing diffusion-weighted imaging of contused spinal cord treated with anti-MBP T cells. Spinal cords of MBP-T cell-treated and PBS-treated animals (with locomotion scores of 10 and 8, respectively) were excised under deep anesthesia, immediately fixed in 4% paraformaldehyde solution, and placed into 5 mm NMR tubes. Diffusion anisotropy was measured in a Bruker DMX 400 widebore spectrometer using a microscopy probe with a 5-mm Helmholtz coil and actively shielded magnetic field gradients. A multislice pulsed gradient spin echo experiment was performed with 9 axial slices, with the central slice positioned at the center of the spinal injury. Images were acquired with TE of 31 ms, TR of 2000 ms, a diffusion time of 15 ms, a diffusion gradient duration of 3 ms, field of view 0.6 mm, matrix size 128 x 128, slice thickness 0.5 mm, and slice separation of 1.18 mm. Four diffusion gradient values of 0, 28, 49, and 71 g/cm were applied along the read direction (transverse diffusion) or along the slice direction (longitudinal diffusion). Diffusion anisotropy is manifested by increased signal intensity in the images with the highest transverse diffusion gradient relative to the longitudinal diffusion gradient. The excised spinal cords of a PBS-treated rat and in the rat treated with MBP-T cells were subjected to diffusion-weighted MRI analysis. In the PBS-treated injured control, diffusion anisotropy was seen mainly in sections near the proximal and distal stumps of the

cord, with low anisotropy in sections taken through the site of injury. In contrast, in the treated rat, higher levels of diffusion anisotropy can be seen in sections taken through the site of injury.

Fig. 10 is a graph illustrating inhibition of secondary degeneration after optic nerve crush injury in adult rats. See text, Section 8, for experimental details. Rats were injected intradermally through the footpads with a 21-mer peptide based on amino acid residues 35-55 (MOG p35-55) of myelin/oligodendrocyte glycoprotein (chemically synthesized at the Weizmann Institute, Israel) ( $50\mu$ /animal) or PBS ten days prior to optic nerve crush injury or MOG p35-55 in the absence of crush injury. MOG p35-55 was administered with Incomplete Freund's Adjuvant. Surviving optic nerve fibers were monitored by retrograde labeling of retinal ganglion cells (RGCs). The number of RGCs in rats injected with PBS or MOG p35-55 was expressed as a percentage of the total number of neurons in rats injected with MOG p35-55 in the absence of crush injury.

Fig. 11 is a graph illustrating inhibition in adult rats of secondary degeneration after optic nerve crush injury by MBP. See text, Section 9, for experimental details. MBP (Sigma, Israel) (1 mg in 0.5 ml saline) was administered orally to adult rats by gavage using a blunt needle. MBP was administered 5 times, i.e., every third day beginning two weeks prior to optic nerve crush injury. Surviving optic nerve fibers were monitored by retrograde labeling of retinal ganglion cells (RGCs). The number of RGCs in treated rats was expressed as a percentage of the total number of neurons in untreated rats following the injury.

Figs. 12 (A-F) show expression of B7 costimulatory molecules in intact and injured rat optic nerve. Optic nerves were excised from adult Lewis rats before (12A, 12B) and three days after injury (12C, 12D, 12E) and analyzed immunohistochemically for expression of the B7 costimulatory molecule. The site of injury was delineated by GFAP staining. Using calibrated cross-action forceps, the right optic nerve was subjected to a mild crush injury 1-2 mm from the eye. The uninjured contralateral nerve was left undisturbed. Immunohistochemical analysis of optic nerve antigens was

performed as follows. Briefly, longitudinal cryosections of the excised nerves (20  $\mu$ m thick) were picked up onto gelatin-coated glass and fixed with ethanol for ten minutes at room temperature. The sections were washed and incubated for one hour at room temperature with mouse monoclonal antibody to rat GFAP (BioMakor, Israel), diluted 1:100, and with antibodies to B7.2 costimulatory molecule and the B7.1 costimulatory molecule (PHARMINGEN, San Diego, CA), diluted 1:25. The sections were washed again and incubated with rhodamine isothiocyanate-conjugated goat anti-mouse IgG (with minimal cross-reaction to rat, human, bovine and horse serum protein) (Jackson ImmunoResearch, West Grove, PA), for one hour at room temperature. All washing solutions contained PBS and 0.05% Tween-20. All diluting solutions contained PBS containing 3% fetal calf serum and 2% bovine serum albumin. The sections were treated with glycerol containing 1,4-diazobicyclo-(2,2,2)-octane and were then viewed with a Zeiss microscope. Note the morphological changes of the B7.2 positive cells after injury, from a rounded (12A, 12B) to a star-like shape (12C, 12D). The B7.2 positive cells were present at a higher density closer to the injury site (12E). Expression of B7.1 was detectable only from day seven and only at the injured site (12F).

Figs. 13 A-C show immunohistochemical analysis of T cells, macrophages or microglia, and B7.2 costimulatory molecules in the injured optic nerves of rats fed MBP. Lewis rats aged 6-8 weeks were fed 1 mg of bovine MBP (Sigma, Israel) (2 mg MBP/ml PBS) or 0.5 ml PBS only every other day by gastric intubation using a stainless steel feeding needle (Thomas Scientific, Swedesboro, NJ) (Chen, Y., Kuchroo, V.K., Inobe, J. Hafler, D.A. & Weiner, H.L. Regulatory T cell clones induced by oral tolerance: suppression of autoimmune encephalomyelitis. Science 265:1237-1240, 1994). Ten days after starting MBP the right optic nerves were subjected to calibrated crush injury, as described for Figure 12. Three days later the nerves were excised and prepared for immunohistochemical analysis of T cells using mouse monoclonal antibodies to T cell receptor 11, diluted 1:25, macrophages or microglia using anti-ED1 antibodies (Serotek, Oxford, U.K) diluted 1:250, astrocytes using anti-GFAP antibodies and B7.2

costimulatory molecules as described for Fig. 12. There were no significant quantitative differences in T cells or in ED-1 positive cells between injured optic nerves of PBS-fed (13A) and MBP-fed (13B) rats. The number of B7.2 positive cells at the site of injury of MBP-fed rats (13C) should be noted, as compared with injured controls (Fig. 12E).

Fig. 14 is a graph showing the slowing of neuronal degeneration in rats with orally induced tolerance to MBP. Lewis rats were fed 1 mg MBP daily, or every other day, or 4 times a day at two hour intervals for five consecutive days. Control animals were given PBS or the non-self antigen OVA (Sigma, Israel). Ten days after the start of MBP ingestion, the right optic nerves were subjected to a calibrated mild crush injury. Two weeks later the RGCs were retrogradely labelled by application of the fluorescent lipophilic dye, 4-(4-didecylamino)styryl)-N-methylpyridinium iodide (4-Di-10-Asp) (Molecular Probes Europe BV, Netherlands), distally to the site of injury, as described. Briefly, complete axotomy was performed 1-2 mm from the distal border to the injury site, and solid crystals (0.2-0.4 mm in diameter) of 4-Di-10-Asp were immediately deposited at the site of the lesion. Retrograde labelling of RGCs by the dye gives a reliable indication of the number of still-functioning neurons, as only intact axons can transport the dye to their cell bodies in the retina. Six days after dye application, the retina was detached from the eye, prepared as a flattened whole mount in 4% paraformaldehyde solution, and examined for labelled ganglion cells by fluorescence microscopy. RGCs were counted from three different regions in the retina. The results are expressed as normalized percentage of each retina to untreated injured animal mean of the same experiment. The median of each group is shown as a bar (Control vs. MBP OTx4 \*\*  $P < 0.01$ ; Control vs. MBP OT \*\*  $P, 0.01$ ; Control vs. OVA OT ns  $P > 0.05$ ).

Fig. 15 shows the nucleotide sequence of rat myelin basic protein gene, SEQ ID NO:1, Genbank accession number M25889 (Schaich et al., Biol. Chem. 367:825-834, 1986).

Fig. 16 shows the nucleotide sequence of human myelin basic protein gene, SEQ ID NO:2, Genbank accession number

M13577 (Kamholz et al., Proc. Natl. Acad. Sci. U.S.A. 83(13): 4962-4966, 1986).

Figs 17 (A-F) show the nucleotide sequences of human myelin proteolipid protein gene exons 1-7, SEQ ID NOs:3-8, respectively, Genbank accession number M15026-M15032 respectively (Diehl et al., Proc. Natl. Acad. Sci. U.S.A. 83(24):9807-9811, 1986; published erratum appears in Proc Natl Acad Sci U.S.A. 86(6):617-8, 1991).

Fig. 18 shows the nucleotide sequence of human myelin oligodendrocyte glycoprotein gene, SEQ ID NO:9, Genbank accession number Z48051 (Roth et al., submitted (17-Jan-1995) Roth, CNRS UPR 8291, CIGH, CHU Purpan, Toulouse, France, 31300; Gonzalez et al., Mol. Phylogent. Evol. 6:63-71, 1996).

Fig. 19 shows the nucleotide sequence of rat proteolipid protein and variant, SEQ ID NO:10, Genbank accession number M16471 (Nave et al, Proc. Natl. Acad. Sci. U.S.A. 84:600-604, 1987).

Fig. 20 shows the nucleotide sequence of rat myelin-associated glycoprotein, SEQ ID NO:11, Genbank accession number M14871 (Arquint et al, Proc. Natl. Acad. Sci. USA 84:600-604, 1987).

Fig. 21 shows the amino acid sequence of human myelin basic protein, SEQ ID NO:12, Genbank accession number 307160 (Kamholz et al., 1986, Proc. Natl. Acad. Sci. U.S.A. 83(13):4962-4966, 1986).

Fig. 22 shows the amino acid sequence of human proteolipid protein, SEQ ID NO:13, Genbank accession number 387028.

Fig. 23 shows the amino acid sequence of human myelin oligodendrocyte glycoprotein, SEQ ID NO:14, Genbank accession number 793839 (Roth et al., Genomics 28(2):241-250, 1995; Roth Submitted (17-JAN-1995) Roth CNRS UPR 8291, CIGH, CHU Purpan, Toulouse, France, 31300; Gonzalez et al., Mol. Phylogent. Evol. 6:63-71, 1996).

#### DETAILED DESCRIPTION OF THE INVENTION

Merely for ease of explanation, the detailed description of the present invention is divided into the following subsections: (1) NS-specific activated T cells; (2)

NS-specific antigens, peptides derived therefrom and derivatives thereof; (3) nucleotide sequences encoding NS-specific antigens and peptides derived therefrom; (4) therapeutic uses of non-recombinant, NS-specific activated T cells, NS-specific antigens, peptides derived therefrom and derivatives thereof, and nucleotide sequences encoding NS-specific antigens and peptides derived therefrom; and (5) formulations and modes of administration of nonrecombinant, NS-specific activated T cells, NS-specific antigens, peptides derived therefrom and derivatives thereof, and nucleotide sequences encoding NS-specific antigens and peptides derived therefrom.

#### 5.1 NS-SPECIFIC ACTIVATED T CELLS

NS-specific activated T cells (ATCs) can be used for ameliorating or inhibiting the effects of injury or disease of the CNS or PNS that result in NS degeneration or for promoting regeneration in the NS, in particular the CNS.

The NS-specific activated T cells are preferably autologous, most preferably of the CD4 and/or CD8 phenotypes, but they may also be allogeneic T cells from related donors, e.g., siblings, parents, children, or HLA-matched or partially matched, semi-allogeneic or fully allogeneic donors.

In addition to the use of autologous T cells isolated from the subject, the present invention also comprehends the use of semi-allogeneic T cells for neuroprotection. These T cells may be prepared as short- or long-term lines and stored by conventional cryopreservation methods for thawing and administration, either immediately or after culturing for 1-3 days, to a subject suffering from injury to the central nervous system and in need of T cell neuroprotection.

The use of semi-allogeneic T cells is based on the fact that T cells can recognize a specific antigen epitope presented by foreign antigen presenting cells (APC), provided that the APC express the MHC molecule, class I or class II, to which the specific responding T cell population is restricted, along with the antigen epitope recognized by the T cells. Thus, a semi-allogeneic population of T cells that can recognize at least one allelic product of the subject's MHC

molecules, preferably an HLA-DR or an HLA-DQ or other HLA molecule, and that is specific for a NS-associated antigen epitope, will be able to recognize the NS antigen in the subject's area of NS damage and produce the needed neuroprotective effect. There is little or no polymorphism in the adhesion molecules, leukocyte migration molecules, and accessory molecules needed for the T cells to migrate to the area of damage, accumulate there, and undergo activation. Thus, the semi-allogeneic T cells will be able to migrate and accumulate at the CNS site in need of neuroprotection and will be activated to produce the desired effect.

It is known that semi-allogeneic T cells will be rejected by the subject's immune system, but that rejection requires about two weeks to develop. Hence, the semi-allogeneic T cells will have the two week window of opportunity needed to exert neuroprotection. After two weeks, the semi-allogeneic T cells will be rejected from the body of the subject, but that rejection is advantageous to the subject because it will rid the subject of the foreign T cells and prevent any untoward consequences of the activated T cells. The semi-allogeneic T cells thus provide an important safety factor and are a preferred embodiment.

It is known that a relatively small number of HLA class II molecules are shared by most individuals in a population. For example, about 50% of the Jewish population express the HLA-DR5 gene. Thus, a bank of specific T cells reactive to NS antigen epitopes that are restricted to HLA-DR5 would be useful in 50% of that population. The entire population can be covered essentially by a small number of additional T cell lines restricted to a few other prevalent HLA molecules, such as DR1, DR4, DR2, etc. Thus, a functional bank of uniform T cell lines can be prepared and stored for immediate use in almost any individual in a given population. Such a bank of T cells would overcome any technical problems in obtaining a sufficient number of specific T cells from the subject in need of neuroprotection during the open window of treatment opportunity. The semi-allogeneic T cells will be safely rejected after accomplishing their role of neuroprotection. This aspect of the invention does not

contradict, and is in addition to the use of autologous T cells as described herein.

The NS-specific activated T cells are preferably non-attenuated, although attenuated NS-specific activated T cells may be used. T cells may be attenuated using methods well known in the art, including but not limited to, by gamma-irradiation, e.g., 1.5-10.0 Rads (Ben-Nun, A., Wekerle, H. and Cohen, I.R., Nature 292:60-61 (1981); Ben-Nun, A. and Cohen, I.R., J. Immunol. 129:303-308 (1982)); and/or by pressure treatment, for example as described in U.S. Patent No. 4,996,194 (Cohen et al.); and/or by chemical cross-linking with an agent such as formaldehyde, glutaraldehyde and the like, for example as described in U.S. Patent No. 4,996,194 (Cohen et al.); and/or by cross-linking and photoactivation with light with a photoactivatable psoralen compound, for example as described in U.S. Patent No. 5,114,721 (Cohen et al.); and/or by a cytoskeletal disrupting agent such as cytochalasin and colchicine, for example as described in U.S. Patent No. 4,996,194 (Cohen et al.). In a preferred embodiment the NS-specific activated T cells are isolated as described below. T cells can be isolated and purified according to methods known in the art (Mor and Cohen, 1995, J. Immunol. 155:3693-3699). For an illustrative example, see Section 6.1.

Circulating T cells of a subject which recognize myelin basic protein or another NS antigen, such as the amyloid precursor protein, are isolated and expanded using known procedures. In order to obtain NS-specific activated T cells, T cells are isolated and the NS-specific ATCs are then expanded by a known procedure (Burns et al., Cell Immunol. 81:435, 1983; Pette et al., Proc. Natl. Acad. Sci. USA 87:7968, 1990; Mortin et al., J. Immunol. 145:540, 1990; Schluesener et al., J. Immunol. 135:3128, 1985; Suruhan-Dires Keneli et al., Euro. J. Immunol. 23:530, 1993, which are incorporated herein by reference in their entirety).

The isolated T cells may be activated by exposure of the cells to one or more of a variety of natural or synthetic NS-specific antigens or epitopes, including but not limited to, myelin basic protein (MBP), myelin oligodendrocyte glycoprotein (MOG), proteolipid protein (PLP), myelin-associated



glycoprotein (MAG), S-100,  $\beta$ -amyloid, Thy-1, P0, P2 and neurotransmitter receptors. In a preferred embodiment, the isolated T cells are activated by one or more cryptic epitopes, including but limited to the following MBP peptides: p11-30, p51-70, p91-110, p131-150, and p-151-170.

During *ex vivo* activation of the T cells, the T cells may be activated by culturing them in medium to which at least one suitable growth promoting factor has been added. Growth promoting factors suitable for this purpose include, without limitation, cytokines, for instance tumor necrosis factor  $\alpha$  (TNF- $\alpha$ ), interleukin 2 (IL-2), and interleukin 4 (IL-4).

In one embodiment, the activated T cells endogenously produce a substance that ameliorates the effects of injury or disease in the NS.

In another embodiment, the activated T cells endogenously produce a substance that stimulates other cells, including, but not limited to, transforming growth factor- $\beta$  (TGF- $\beta$ ), nerve growth factor (NGF), neurotrophic factor 3 (NT-3), neurotrophic factor 4/5 (NT-4/5), brain derived neurotrophic factor (BDNF); interferon- $\gamma$  (IFN- $\gamma$ ), and interleukin-6 (IL-6), wherein the other cells, directly or indirectly, ameliorate the effects of injury or disease.

Following their proliferation *in vitro*, the T cells are administered to a mammalian subject. In a preferred embodiment, the T cells are administered to a human subject. T cell expansion is preferably performed using peptides corresponding to sequences in a non-pathogenic, NS-specific, self protein.

A subject can initially be immunized with an NS-specific antigen using a non-pathogenic peptide of the self protein. A T cell preparation can be prepared from the blood of such immunized subjects, preferably from T cells selected for their specificity towards the NS-specific antigen. The selected T cells can then be stimulated to produce a T cell line specific to the self-antigen (Ben-Nun et al., J. Immunol. 129:303, 1982).

The NS-specific antigen may be a purified antigen or a crude NS preparation, as will be described below. NS-

specific antigen activated T cells, obtained as described above, can be used immediately or may be preserved for later use, e.g., by cryopreservation as described below. NS-specific activated T cells may also be obtained using previously cryopreserved T cells, i.e., after thawing the cells, the T cells may be incubated with NS-specific antigen, optimally together with thymocytes, to obtain a preparation of NS-specific ATCs.

As will be evident to those skilled in the art, the T cells can be preserved, e.g., by cryopreservation, either before or after culture.

Cryopreservation agents which can be used include but are not limited to dimethyl sulfoxide (DMSO) (Lovelock and Bishop, Nature 183:1394-1395, 1959; Ashwood-Smith, Nature 190:1204-1205, 1961), glycerol, polyvinylpyrrolidone (Rinfret, Ann. N.Y. Acad. Sci. 85:576, 1960), polyethylene glycol (Sloviter and Ravdin, Nature 196:548, 1962), albumin, dextran, sucrose, ethylene glycol, i-erythritol, D-ribitol, D-mannitol (Rowe et al., Fed. Proc. 21:157, 1962), D-sorbitol, i-inositol, D-lactose, choline chloride (Bender et al., J. Appl. Physiol. 15:520, 1960), amino acids (Phan The Tran and Bender, Exp. Cell Res. 20:651, 1960), methanol, acetamide, glycerol monoacetate (Lovelock, Biochem. J. 56:265, 1954), inorganic salts (Phan The Tran and Bender, Proc. Soc. Exp. Biol. Med. 104:388, 1960; Phan The Tran and Bender, 1961, in Radiobiology, Proceedings of the Third Australian Conference on Radiobiology, Ilbery, P.L.T., ed., Butterworth, London, p. 59), and DMSO combined with hydroxyethyl starch and human serum albumin (Zaroulis and Leiderman, Cryobiology 17:311-317, 1980).

A controlled cooling rate is critical. Different cryoprotective agents (Rapatz et al., Cryobiology 5(1):18-25, 1968) and different cell types have different optimal cooling rates. See, e.g., Rowe and Rinfret, Blood 20:636 (1962); Rowe, Cryobiology 3(1):12-18 (1966); Lewis et al., Transfusion 7(1):17-32 (1967); and Mazur, Science 168:939-949 (1970) for effects of cooling velocity on survival of cells and on their transplantation potential. The heat of fusion phase where water turns to ice should be minimal. The cooling procedure

can be carried out by use of, e.g., a programmable freezing device or a methanol bath procedure.

Programmable freezing apparatuses allow determination of optimal cooling rates and facilitate standard reproducible cooling. Programmable controlled-rate freezers such as Cryomed or Planar permit tuning of the freezing regimen to the desired cooling rate curve.

After thorough freezing, cells can be rapidly transferred to a long-term cryogenic storage vessel. In one embodiment, samples can be cryogenically stored in mechanical freezers, such as freezers that maintain a temperature of about  $-80^{\circ}\text{C}$  or about  $-20^{\circ}\text{C}$ . In a preferred embodiment, samples can be cryogenically stored in liquid nitrogen ( $-196^{\circ}\text{C}$ ) or its vapor. Such storage is greatly facilitated by the availability of highly efficient liquid nitrogen refrigerators, which resemble large Thermos containers with an extremely low vacuum and internal super insulation, such that heat leakage and nitrogen losses are kept to an absolute minimum.

Considerations and procedures for the manipulation, cryopreservation, and long term storage of T cells can be found, for example, in the following references, incorporated by reference herein: Gorin, Clinics in Haematology 15(1):19-48 (1986); Bone-Marrow Conservation, Culture and Transplantation, Proceedings of a Panel, Moscow, July 22-26, 1968, International Atomic Energy Agency, Vienna, pp. 107-186.

Other methods of cryopreservation of viable cells, or modifications thereof, are available and envisioned for use, e.g., cold metal-mirror techniques. See Livesey and Linner, Nature 327:255 (1987); Linner et al., J. Histochem. Cytochem. 34(9):1123-1135 (1986); see also U.S. Patent No. 4,199,022 by Senken et al., U.S. Patent No. 3,753,357 by Schwartz, U.S. Patent No. 4,559,298 by Fahy.

Frozen cells are preferably thawed quickly (e.g., in a water bath maintained at  $37-47^{\circ}\text{C}$ ) and chilled immediately upon thawing. It may be desirable to treat the cells in order to prevent cellular clumping upon thawing. To prevent clumping, various procedures can be used, including but not limited to the addition before or after freezing of DNase (Spitzer et al., Cancer 45:3075-3085, 1980), low molecular

weight dextran and citrate, citrate, hydroxyethyl starch (Stiff et al., Cryobiology 20:17-24, 1983), or acid citrate dextrose (Zaroulis and Leiderman, Cryobiology 17:311-317, 1980), etc.

The cryoprotective agent, if toxic in humans, should be removed prior to therapeutic use of the thawed T cells. One way in which to remove the cryoprotective agent is by dilution to an insignificant concentration.

Once frozen T cells have been thawed and recovered, they are used to promote neuronal regeneration as described herein with respect to non-frozen T cells. Once thawed, the T cells may be used immediately, assuming that they were activated prior to freezing. Preferably, however, the thawed cells are cultured before injection to the patient in order to eliminate non-viable cells. Furthermore, in the course of this culturing over a period of about one to three days, an appropriate activating agent can be added so as to activate the cells, if the frozen cells were resting T cells, or to help the cells achieve a higher rate of activation if they were activated prior to freezing. Usually, time is available to allow such a culturing step prior to administration as the T cells may be administered as long as a week after injury, and possibly longer, and still maintain their neuroregenerative and neuroprotective effect.

## 5.2 NS-SPECIFIC ANTIGENS AND PEPTIDES DERIVED THEREFROM

Pharmaceutical compositions comprising an NS-specific antigen or peptide derived therefrom or derivative thereof can be used for preventing or inhibiting the effects of injury or disease that result in NS degeneration or for promoting nerve regeneration in the NS, particularly in the CNS. Additionally, NS-specific antigens or peptides derived therefrom or derivatives thereof may be used for in vivo or in vitro activation of T cells. In one embodiment, the NS-specific antigen is an isolated or purified antigen. In another embodiment, methods of promoting nerve regeneration or of preventing or inhibiting the effects of CNS or PNS injury or disease comprise administering NS-specific antigen or a peptide derived therefrom or derivative thereof to a mammal wherein the

NS-specific antigen or peptide derived therefrom or derivative thereof activates T cells *in vivo* to produce a population of T cells that accumulate at a site of injury or disease of the CNS or PNS.

The NS-specific antigen may be an antigen obtained from NS tissue, preferably from tissue at a site of CNS injury or disease. The NS-specific antigen may be isolated and purified by standard methods including chromatography (e.g., ion exchange, affinity, and sizing column chromatography), centrifugation, differential solubility, or by any other standard technique for the purification of antigens. The functional properties may be evaluated using any suitable assay. In the practice of the invention, natural or synthetic NS-specific antigens or epitopes include, but are not limited to, MBP, MOG, PLP, MAG, S-100,  $\beta$ -amyloid, Thy-1, P0, P2 and a neurotransmitter receptor.

Specific illustrative examples of useful NS-specific antigens include but are not limited to, human MBP, depicted in Fig. 21, (SEQ ID NO:12); human proteolipid protein, depicted in Fig. 22 (SEQ ID NO:13); and human oligodendrocyte glycoprotein, depicted in Fig. 23 (SEQ ID NO:14).

In a preferred embodiment, peptides derived from NS-specific, self-antigens or derivatives of NS-specific antigens activate T cells, but do not induce an autoimmune disease. An example of such peptide is a peptide comprising amino acids 51-70 of myelin basic protein (residues 51-70 of SEQ ID NO:12).

In addition, an NS-specific antigen may be a crude NS-tissue preparation, e.g., derived from NS tissue obtained from mammalian NS. Such a preparation may include cells, both living or dead cells, membrane fractions of such cells or tissue, etc.

an NS-specific antigen may be obtained by an NS biopsy or necropsy from a mammal including, but not limited to, from a site of CNS injury; from cadavers; from cell lines grown in culture. Additionally, an NS-specific antigen may be a protein obtained by genetic engineering, chemically synthesized, etc.

In addition to NS-specific antigens, the invention also relates to peptides derived from NS-specific antigens or

derivatives including chemical derivatives and analogs of NS-specific antigens which are functionally active, i.e., they are capable of displaying one or more known functional activities associated with a full-length NS-specific antigen. Such functional activities include but are not limited to antigenicity (ability to bind (or compete with an NS-antigen for binding) to an anti-NS-specific antibody), immunogenicity (ability to generate antibody which binds to an NS-specific protein), and ability to interact with T cells, resulting in activation comparable to that obtained using the corresponding full-length antigen. The crucial test is that the antigen which is used for activating the T cells causes the T cells to be capable of recognizing an antigen in the NS of the mammal (patient) being treated.

A peptide derived from a CNS-specific or PNS-specific antigen preferably has a sequence comprised within the antigen sequence and is either: (1) an immunogenic peptide, i.e., a peptide that can elicit a human T cell response detected by a T cell proliferation or by cytokine (e.g. interferon (IFN)- $\gamma$ , interleukin (IL)-2, IL-4 or IL-10) production or (2) a "cryptic epitope" (also designated herein as "immunosilent" or "nonimmunodominant" epitope), i.e., a peptide that by itself can induce a T cell immune response that is not induced by the whole antigen protein (see Moalem et al., Nature Med. 5(1), 1999). Cryptic epitopes for use in the present invention include, but are not limited to, peptides of the myelin basic protein sequence: peptide p11-30, p51-70, p91-110, p131-150, and p151-170. Other peptides can be identified by their capacity to elicit a human T cell response detected by T cell proliferation or by cytokine (e.g. IFN- $\gamma$ , IL-2, IL-4, or IL-10) production. Such cryptic epitopes are particularly preferred as T cells activated thereby will accumulate at the injury site, in accordance with the present invention, but are particularly weak in autoimmunity. Thus, they would be expected to have fewer side effects.

In one specific embodiment of the invention, peptides consisting of or comprising a fragment of an NS-specific antigen consisting of at least 10 (contiguous) amino acids of the NS-specific antigen are provided. In other embodiments, the

fragment consists of at least 20 contiguous amino acids or 50 contiguous amino acids of the NS-specific antigen. Derivatives of an NS-specific antigen also include but are not limited to those molecules comprising regions that are substantially homologous to the full-length antigen or fragments thereof (e.g., in various embodiments, at least 60% or 70% or 80% or 90% or 95% identity over an amino acid sequence of identical size or when compared to an aligned sequence in which the alignment is done by a computer homology program known in the art) or whose encoding nucleic acid is capable of hybridizing to a coding nucleotide sequence of the full-length NS-specific antigen, under high stringency, moderate stringency, or low stringency conditions.

Computer programs for determining homology may include but are not limited to TBLASTN, BLASTP, FASTA, TFASTA, and CLUSTALW (Pearson and Lipman, Proc. Natl. Acad. Sci. USA 85(8):2444-8, 1988; Altschul et al., J. Mol. Biol. 215(3):40310, 1990; Thompson, et al., Nucleic Acids Res. 22(22):4673-80, 1994; Higgins, et al., Methods Enzymol 266:383-402, 1996; Altschul, et al., 1990, J. Mol. Biol. 215(3):403-410, 1990).

The NS-specific antigen derivatives of the invention can be produced by various methods known in the art. The manipulations which result in their production can occur at the gene or protein level. For example, a cloned gene sequence can be modified by any of numerous strategies known in the art (Maniatis, T., 1990, Molecular Cloning, A Laboratory Manual, 2d ed., Cold Spring Harbor Laboratory, Cold Spring Harbor, New York). The sequence can be cleaved at appropriate sites with restriction endonuclease(s), followed by further enzymatic modification if desired, isolated, and ligated in vitro.

Additionally, the coding nucleic acid sequence can be mutated in vitro or in vivo, to create and/or destroy translation, initiation, and/or termination sequences, or to create variations in coding regions and/or form new restriction endonuclease sites or destroy preexisting ones, to facilitate further in vitro modification. Any technique for mutagenesis known in the art can be used, including but not limited to,

chemical mutagenesis, *in vitro* site-directed mutagenesis (Hutchinson, C., et al., J. Biol. Chem 253:6551, 1978), etc.

Manipulations may also be made at the protein level. Included within the scope of the invention are derivatives which are differentially modified during or after translation, e.g., by glycosylation, acetylation, phosphorylation, amidation, derivatization by known protecting/blocking groups, proteolytic cleavage, linkage to an antibody molecule or other cellular ligand, etc. Any of numerous chemical modifications may be carried out by known techniques, including but not limited to specific chemical cleavage by cyanogen bromide, trypsin, chymotrypsin, papain, V8 protease,  $\text{NaBH}_4$ ; acetylation, formylation, oxidation, reduction; metabolic synthesis in the presence of tunicamycin; etc.

In addition, derivatives of an NS-specific antigen can be chemically synthesized. For example, a peptide corresponding to a portion of an antigen which comprises the desired domain or which mediates the desired activity can be synthesized by use of a peptide synthesizer. Furthermore, if desired, nonclassical amino acids or chemical amino acids analogs can be introduced as a substitution or addition into the amino acid sequence. Non-classical amino acids include but are not limited to the D-isomers of the common amino acids,  $\alpha$ -amino isobutyric acid; 4-aminobutyric acid, Abu; 2-amino butyric acid,  $\gamma$ -Abu;  $\epsilon$ -Ahx, 6-amino hexanoic acid; Aib, 2-amino isobutyric acid; 3-amino propionic acid; ornithine; norleucine; novaline; hydroxyproline; sarcosine; citrulline; cysteic acid; t-butylglycine; t-butylalanine; phenylglycine; cyclohexylalanine;  $\beta$ -alanine; fluoro-amino acids; designer amino acids such as  $\beta$ -methyl amino acids,  $\text{C}\alpha$ -methyl amino acids,  $\text{N}\alpha$ -methyl amino acids, and amino acid analogs in general. Furthermore, the amino acid can be D (dextrorotary) or L (levorotary).

The functional activity of NS-specific antigens and peptides derived therefrom and derivatives thereof can be assayed by various methods known in the art, including, but not limited to, T cell proliferation assays (Mor and Cohen, J. Immunol. 155:3693-3699, 1995).



An NS-specific antigen or peptide derived therefrom or derivative thereof may be kept in solution or may be provided in a dry form, e.g. as a powder or lyophilizate, to be mixed with appropriate solution prior to use.

### 5.3 NUCLEOTIDE SEQUENCES ENCODING NS-ANTIGENS AND PEPTIDES DERIVED THEREFROM

Compositions comprising a nucleotide sequence encoding an NS-specific antigen or peptide derived therefrom can be used for preventing or inhibiting the effects of injury or disease that result in CNS or PNS degeneration or for promoting nerve regeneration in the CNS or PNS. Specific illustrative examples of useful nucleotide sequences encoding NS-specific antigens or peptides derived from an NS-specific antigen, include but are not limited to nucleotide sequences encoding rat myelin basic protein (MBP) peptides, depicted in Fig. 15 (SEQ ID NO:1); human MBP, depicted in Fig. 16 (SEQ ID NO:2); human myelin PLP, depicted in Figs. 17(A-F) (SEQ ID NOs:3-8); human MOG, depicted in Fig. 18 (SEQ ID NO:9); rat PLP and variant, depicted in Fig. 19 (SEQ ID NO:10); and rat MAG, depicted in Fig. 20 (SEQ ID NO:11).

### 5.4 THERAPEUTIC USES

The compositions described in Sections 5.1 through 5.3 may be used to promote nerve regeneration or to prevent or inhibit secondary degeneration which may otherwise follow primary NS injury, e.g., blunt trauma, penetrating trauma, hemorrhagic stroke, ischemic stroke or damages caused by surgery such as tumor excision. In addition, such compositions may be used to ameliorate the effects of disease that result in a degenerative process, e.g., degeneration occurring in either grey or white matter (or both) as a result of various diseases or disorders, including, without limitation: diabetic neuropathy, senile dementias, Alzheimer's disease, Parkinson's Disease, facial nerve (Bell's) palsy, glaucoma, Huntington's chorea, amyotrophic lateral sclerosis (ALS), non-arteritic optic neuropathy, intervertebral disc herniation, vitamin deficiency, prion diseases such as Creutzfeldt-Jakob disease, carpal tunnel syndrome, peripheral neuropathies associated with

various diseases, including but not limited to, uremia, porphyria, hypoglycemia, Sjorgren Larsson syndrome, acute sensory neuropathy, chronic ataxic neuropathy, biliary cirrhosis, primary amyloidosis, obstructive lung diseases, acromegaly, malabsorption syndromes, polycythemia vera, IgA and IgG gammopathies, complications of various drugs (e.g., metronidazole) and toxins (e.g., alcohol or organophosphates), Charcot-Marie-Tooth disease, ataxia telangiectasia, Friedreich's ataxia, amyloid polyneuropathies, adrenomyeloneuropathy, Giant axonal neuropathy, Refsum's disease, Fabry's disease, lipoproteinemia, etc.

In a preferred embodiment, the NS-specific activated T cells, the NS-specific antigens, peptides derived therefrom, derivatives thereof or the nucleotides encoding said antigens, or peptides or any combination thereof of the present invention are used to treat diseases or disorders where promotion of nerve regeneration or prevention or inhibition of secondary neural degeneration is indicated, which are not autoimmune diseases or neoplasias. In a preferred embodiment, the compositions of the present invention are administered to a human subject.

While activated NS-specific T cells may have been used in the prior art in the course of treatment to develop tolerance to autoimmune antigens in the treatment of autoimmune diseases, or in the course of immunotherapy in the treatment of NS neoplasms, the present invention can also be used to ameliorate the degenerative process caused by autoimmune diseases or neoplasms as long as it is used in a manner not suggested by such prior art methods. Thus, for example, T cells activated by an autoimmune antigen have been suggested for use to create tolerance to the autoimmune antigen and, thus, ameliorate the autoimmune disease. Such treatment, however, would not have suggested the use of T cells directed to other NS antigens or NS antigens which will not induce tolerance to the autoimmune antigen or T cells which are administered in such a way as to avoid creation of tolerance. Similarly, for neoplasms, the effects of the present invention can be obtained without using immunotherapy processes suggested in the prior art by, for example, using an NS antigen which

does not appear in the neoplasm. T cells activated with such an antigen will still accumulate at the site of neural degeneration and facilitate inhibition of this degeneration, even though it will not serve as immunotherapy for the tumor *per se*.

### 5.5 FORMULATIONS AND ADMINISTRATION

Pharmaceutical compositions for use in accordance with the present invention may be formulated in conventional manner using one or more physiologically acceptable carriers or excipients. The carrier(s) must be "acceptable" in the sense of being compatible with the other ingredients of the composition and not deleterious to the recipient thereof.

The term "carrier" refers to a diluent, adjuvant, excipient, or vehicle with which the therapeutic is administered. The carriers in the pharmaceutical composition may comprise a binder, such as microcrystalline cellulose, polyvinylpyrrolidone (polyvidone or povidone), gum tragacanth, gelatin, starch, lactose or lactose monohydrate; a disintegrating agent, such as alginic acid, maize starch and the like; a lubricant or surfactant, such as magnesium stearate, or sodium lauryl sulphate; a glidant, such as colloidal silicon dioxide; a sweetening agent, such as sucrose or saccharin; and/or a flavoring agent, such as peppermint, methyl salicylate, or orange flavoring.

Methods of administration include, but are not limited to, parenteral, e.g., intravenous, intraperitoneal, intramuscular, subcutaneous, mucosal (e.g., oral, intranasal, buccal, vaginal, rectal, intraocular), intrathecal, topical and intradermal routes. Administration can be systemic or local.

For oral administration, the pharmaceutical preparation may be in liquid form, for example, solutions, syrups or suspensions, or may be presented as a drug product for reconstitution with water or other suitable vehicle before use. Such liquid preparations may be prepared by conventional means with pharmaceutically acceptable additives such as suspending agents (e.g., sorbitol syrup, cellulose derivatives or hydrogenated edible fats); emulsifying agents (e.g., lecithin or acacia); non-aqueous vehicles (e.g., almond oil,

oily esters, or fractionated vegetable oils); and preservatives (e.g., methyl or propyl-p-hydroxybenzoates or sorbic acid). The pharmaceutical compositions may take the form of, for example, tablets or capsules prepared by conventional means with pharmaceutically acceptable excipients such as binding agents (e.g., pregelatinized maize starch, polyvinyl pyrrolidone or hydroxypropyl methylcellulose); fillers (e.g., lactose, microcrystalline cellulose or calcium hydrogen phosphate); lubricants (e.g., magnesium stearate, talc or silica); disintegrants (e.g., potato starch or sodium starch glycolate); or wetting agents (e.g., sodium lauryl sulphate). The tablets may be coated by methods well-known in the art.

Preparations for oral administration may be suitably formulated to give controlled release of the active compound.

For buccal administration, the compositions may take the form of tablets or lozenges formulated in conventional manner.

The compositions may be formulated for parenteral administration by injection, e.g., by bolus injection or continuous infusion. Formulations for injection may be presented in unit dosage form, e.g., in ampoules or in multidose containers, with an added preservative. The compositions may take such forms as suspensions, solutions or emulsions in oily or aqueous vehicles, and may contain formulatory agents such as suspending, stabilizing and/or dispersing agents. Alternatively, the active ingredient may be in powder form for constitution with a suitable vehicle, e.g., sterile pyrogen free water, before use.

The compositions may also be formulated in rectal compositions such as suppositories or retention enemas, e.g., containing conventional suppository bases such as cocoa butter or other glycerides.

For administration by inhalation, the compositions for use according to the present invention are conveniently delivered in the form of an aerosol spray presentation from pressurized packs or a nebulizer, with the use of a suitable propellant, e.g., dichlorodifluoromethane, trichlorofluoromethane, dichlorotetrafluoroethane, carbon dioxide or other suitable gas. In the case of a pressurized

aerosol the dosage unit may be determined by providing a valve to deliver a metered amount. Capsules and cartridges of, e.g., gelatin, for use in an inhaler or insufflator may be formulated containing a powder mix of the compound and a suitable powder base such as lactose or starch.

In a preferred embodiment, compositions comprising NS-specific activated T cells, an NS-specific antigen or peptide derived therefrom, or derivative thereof, or a nucleotide sequence encoding such antigen or peptide, are formulated in accordance with routine procedures as pharmaceutical compositions adapted for intravenous or intraperitoneal administration to human beings. Typically, compositions for intravenous administration are solutions in sterile isotonic aqueous buffer. Where necessary, the composition may also include a solubilizing agent and a local anesthetic such as lignocaine to ease pain at the site of the injection. Generally, the ingredients are supplied either separately or mixed together. Where the composition is to be administered by infusion, it can be dispensed with an infusion bottle containing sterile pharmaceutical grade water or saline. Where the composition is administered by injection, an ampoule of sterile water or saline for injection can be provided so that the ingredients may be mixed prior to administration.

Pharmaceutical compositions comprising NS-specific antigen or peptide derived therefrom or derivative thereof may optionally be administered with an adjuvant, such as Incomplete Freund's Adjuvant.

The invention also provides a pharmaceutical pack or kit comprising one or more containers filled with one or more of the ingredients of the pharmaceutical compositions of the invention.

In a preferred embodiment, the pharmaceutical compositions of the invention are administered to a mammal, preferably a human, shortly after injury or detection of a degenerative lesion in the NS. The therapeutic methods of the invention may comprise administration of an NS-specific activated T cell or an NS-specific antigen or peptide derived therefrom or derivative thereof, or a nucleotide sequence encoding such antigen or peptide, or any combination thereof.

When using combination therapy, the NS-specific antigen may be administered before, concurrently or after administration of NS-specific activated T cells, a peptide derived from an NS-specific antigen or derivative thereof or a nucleotide sequence encoding such antigen or peptide.

In one embodiment, the compositions of the invention are administered in combination with one or more of the following (a) mononuclear phagocytes, preferably cultured monocytes (as described in PCT publication No. WO 97/09985, which is incorporated herein by reference in its entirety), that have been stimulated to enhance their capacity to promote neuronal regeneration; (b) a neurotrophic factor such as acidic fibroblast growth factor; and (c) an anti-inflammatory therapeutic substance (i.e., an anti-inflammatory steroid, such as dexamethasone or methylprednisolone, or a non-steroidal anti-inflammatory peptide, such as Thr-Lys-Pro (TKP)).

In another embodiment, mononuclear phagocyte cells according to PCT Publication No. WO 97/09985 and U.S. patent application Serial No. 09/041,280, filed March 11, 1998, are injected into the site of injury or lesion within the CNS, either concurrently, prior to, or following parenteral administration of NS-specific activated T cells, an NS-specific antigen or peptide derived therefrom or derivative thereof, or a nucleotide sequence encoding such antigen or peptide.

In another embodiment, administration of NS-specific activated T cells, NS-specific antigen or peptide sequence encoding such antigen or peptide, may be administered as a single dose or may be repeated, preferably at 2 week intervals and then at successively longer intervals once a month, once a quarter, once every six months, etc. The course of treatment may last several months, several years or occasionally also through the life-time of the individual, depending on the condition or disease which is being treated. In the case of a CNS injury, the treatment may range between several days to months or even years, until the condition has stabilized and there is no or only a limited risk of development of secondary degeneration. In chronic human disease or Parkinson's disease, the therapeutic treatment in accordance with the invention may be for life.

As will be evident to those skilled in the art, the therapeutic effect depends at times on the condition or disease to be treated, on the individual's age and health condition, on other physical parameters (e.g. gender, weight, etc.) of the individual, as well as on various other factors, e.g., whether the individual is taking other drugs, etc.

The optimal dose of the therapeutic compositions comprising NS-specific activated T cells of the invention is proportional to the number of nerve fibers affected by NS injury or disease at the site being treated. In a preferred embodiment, the dose ranges from about  $5 \times 10^6$  to about  $10^7$  for treating a lesion affecting about  $10^5$  nerve fibers, such as a complete transection of a rat optic nerve, and ranges from about  $10^7$  to about  $10^8$  for treating a lesion affecting about  $10^6$  -  $10^7$  nerve fibers, such as a complete transection of a human optic nerve. As will be evident to those skilled in the art, the dose of T cells can be scaled up or down in proportion to the number of nerve fibers thought to be affected at the lesion or site of injury being treated.

#### 5.6 ESTABLISHMENT OF AUTOLOGOUS CELL BANKS FOR T LYMPHOCYTES

To minimize secondary damage after nerve injury, patients can be treated by administering autologous or semi-allogeneic T lymphocytes sensitized to at least one appropriate NS antigen. As the window of opportunity has not yet been precisely defined, therapy should be administered as soon as possible after the primary injury to maximize the chances of success, preferably within about one week.

To bridge the gap between the time required for activation and the time needed for treatment, a bank can be established with personal vaults of autologous T lymphocytes prepared for future use for neuroprotective therapy against secondary degeneration in case of NS injury. T lymphocytes are isolated from the blood and then sensitized to a NS antigen. The cells are then frozen and suitably stored under the person's name, identity number, and blood group, in a cell bank until needed.

Additionally, autologous stem cells of the CNS can be processed and stored for potential use by an individual patient in the event of traumatic disorders of the NS such as ischemia or mechanical injury, as well as for treated neurodegenerative conditions such as Alzheimer's disease or Parkinson's disease. Alternatively, semi-allogeneic or allogeneic T cells can be stored frozen in banks for use by any individual who shares one MHC type II molecule with the source of the T cells.

The following examples illustrate certain features of the present invention but are not intended to limit the scope of the present invention.

**EXAMPLE: ACCUMULATION OF ACTIVATED T CELLS IN INJURED  
OPTIC NERVE**

**6.1 MATERIALS AND METHODS**

**6.1.1 ANIMALS**

Female Lewis rats were supplied by the Animal Breeding Center of the Weizmann Institute of Science (Rehovot, IL), matched for age (8-12 weeks) and housed four to a cage in a light and temperature-controlled room.

**6.1.2 MEDIA**

The T cell proliferation medium contained the following: Dulbecco's modified Eagle's medium (DMEM, Biological 15 Industries, Israel) supplemented with 2mM L-glutamine (L-Glu, Sigma, USA),  $5 \times 10^{-5}$  M 2-mercaptoethanol (2-ME, Sigma), penicillin (100 IU/ml; Biological Industries), streptomycin (100  $\mu$ /ml; Biological Industries), sodium pyruvate (1 mM; Biological Industries), non-essential amino acids (1 ml/100 ml; Biological Industries) and autologous rat serum 1% (vol/vol) (Mor et al., Clin. Invest. 85:1594, 1990). Propagation medium contained: DMEM, 2-ME, L-Glu, sodium pyruvate, non-essential amino acids and antibiotics in the same concentration as above with the addition of 10% fetal calf serum (FCS), and 10% T cell growth factor (TCGF) obtained from the supernatant of concanavalin A-stimulated spleen cells (Mor et al., *supra*, 1990).



#### 6.1.3 ANTIGENS

Myelin basic protein (MBP) from the spinal cords of guinea pigs was prepared as described (Hirshfeld, et al., FEBS Lett. 7:317, 1970). Ovalbumin was purchased from Sigma (St. Louis, Missouri). The p51-70 of the rat 18.5kDa isoform of MBP (sequence: APKRGSGKDSHTRTTHYG) (SEQ ID NO:15) and the p277 peptide of the human hsp60 (sequence: VLGGGCALLRCPALDSLTPANED) (SEQ ID NO:16) (Elias et al., Proc. Natl. Acad. Sci. USA 88:3088-3091, 1991) were synthesized using the 9-fluorenylmethoxycarbonyl technique with an automatic multiple peptide synthesizer (AMS 422, ABIMED, Langenfeld, Germany). The purity of the peptides was analyzed by HPLC and amino acid composition.

#### 6.1.4 T CELL LINES

T cell lines were generated from draining lymph node cells obtained from Lewis rats immunized with an antigen (described above in Section 6.1.3). The antigen was dissolved in PBS (1mg/ml) and emulsified with an equal volume of incomplete Freund's adjuvant (Difco Laboratories, Detroit, Michigan) supplemented with 4 mg/ml *Mycobacterium tuberculosis* (Difco 15 Laboratories, Detroit, Michigan). The emulsion (0.1 ml) was injected into hind foot pads of the rats. Ten days after the antigen was injected, the rats were killed and draining lymph nodes were surgically removed and dissociated. The cells were washed and activated with the antigen (10 µg/ml) in proliferation medium (described above in Section 6.1.2). After incubation for 72 h at 37°C, 90% relative humidity and 7% CO<sub>2</sub>, the cells were transferred to propagation medium (described above in Section 6.1.2). Cells were grown in propagation medium for 4-10 days before being re-exposed to antigen (10 µg/ml) in the presence of irradiated (2000 red) thymus cells (10<sup>7</sup> cells/ml) in proliferation medium. The T cell lines were expanded by repeated re-exposure and propagation.

#### 6.1.5 CRUSH INJURY OF RAT OPTIC NERVE

Crush injury of the optic nerve was performed as

previously described (Duvdevani et al., Neurol. Neurosci. 2:31-38, 1990). Briefly, rats were deeply anesthetized by i.p. injection of Rompum (xylazine, 10 mg/kg; Vitamed, Israel) and Vetalar (ketamine, 50 mg/kg; Fort Dodge Laboratories, Fort Dodge, Iowa). Using a binocular operating microscope, a lateral canthotomy was performed in the right eye and the conjunctiva was incised lateral to the cornea. After separation of the retractor bulbi muscles, the optic nerve was exposed intraorbitally by blunt dissection. Using calibrated cross-action forceps, a moderate crush injury was inflicted on the optic nerve, 2mm from the eye (Duvdevani et al., Instructure Neurology and Neuroscience 2:31, 1990). The contralateral nerve was left undisturbed and was used as a control.

#### 6.1.6 IMMUNOCYTOCHEMISTRY OF T CELLS

Longitudinal cryostat nerve sections (20  $\mu$ m thick) were picked up onto gelatin glass slides and frozen until preparation for fluorescent staining. Sections were thawed and fixed in ethanol for 10 minutes at room temperature, washed twice with double-distilled water (ddH<sub>2</sub>O), and incubated for 3 minutes in PBS containing 0.05% polyoxyethylene-sorbitan monolaurate (Tween-20; Sigma, USA). Sections were then incubated for 1 hr at room temperature with a mouse monoclonal antibody directed against rat T cell receptor (TCR) (1:100, Hunig et al., J. Exp. Med., 169:73, 1989), in PBS containing 3% FCS and 2% BSA. After three washes with PBS containing 0.05% Tween-20, the sections were incubated with fluorescein isothiocyanate-conjugated goat anti-mouse IgG (with minimal cross-section to rat, human, bovine and horse serum proteins) (Jackson ImmunoResearch, West Grove, Pennsylvania) for one hour at room temperature. The sections were then washed with PBS containing Tween-20 and treated with glycerol containing 1,4-diazobicyclo-(2,2,2) octane (Sigma), to inhibit quenching of fluorescence. The sections were viewed with a Zeiss microscope and cells were counted. Staining in the absence of first antibody was negative.

## 6.2. RESULTS

Fig. 1 shows accumulation of T cells measured immunohistochemically. The number of T cells was considerably higher in injured nerves rats injected with anti-MBP, anti-OVA or anti-p277 cells; statistical analysis (one-way ANOVA) showed significant differences between T cell numbers in injured optic nerves of rats injected with anti-MBP, anti-OVA, or anti-p277 T cells and in injured optic nerves of rats injected with PBS ( $P < 0.001$ ); and between injured optic nerves and uninjured optic nerves of rats injected with anti-MBP, anti-OVA, or anti-p277 T cells ( $P < 0.001$ ).

### EXAMPLE: NEURPROTECTION BY AUTOIMMUNE ANTI-MBP T CELLS

#### 7.1 MATERIAL AND METHODS

Animals, media, antigens, crush injury of rat optic nerve, sectioning of nerves, T cell lines, and immunolabeling of nerve sections are described in Section 6, *supra*.

##### 7.1.1. RETROGRADE LABELING AND MEASUREMENT OF PRIMARY DAMAGE AND SECONDARY DEGENERATION

Primary damage of the optic nerve axons and their attached retinal ganglion cells (RGCs) were measured after the immediate post-injury application of the fluorescent lipophilic dye 4-(4-(didecylamino)styryl)-N-methylpyridinium iodide (4-Di-Asp) (Molecular Probes Europe BV, Netherlands) distal to the site of injury. Only axons that are intact are capable of transporting the dye back to their cell bodies; therefore, the number of labeled cell bodies is a measure of the number of axons that survived the primary damage. Secondary degeneration was also measured by application of the dye distal to the injury site, but two weeks after the primary lesion was inflicted. Application of the neurotracer dye distal to the site of the primary crush after two weeks ensures that only axons that survived both the primary damage and the secondary degeneration will be counted. This approach makes it possible to differentiate between neurons that are still functionally intact and neurons in which the axons are injured but the cell

bodies are still viable, as only those neurons whose fibers are morphologically intact can take up dye applied distally to the site of injury and transport it to their cell bodies. Using this method, the number of labeled ganglion cells reliably reflects the number of still-functioning neurons. Labeling and measurement were done by exposing the right optic nerve for a second time, again without damaging the retinal blood supply. Complete axotomy was done 1-2 mm from the distal border of the injury site and solid crystals (0.2-0.4 mm in diameter) of 4-Di-10-Asp were deposited at the site of the newly formed axotomy. Uninjured optic nerves were similarly labeled at approximately the same distance from the globe. Five days after dye application, the rats were killed. The retina was detached from the eye, prepared as a flattened whole mount in 4% paraformaldehyde solution and examined for labeled ganglion cells by fluorescence microscopy. The percentage of RGCs surviving secondary degeneration was calculated using the following formula: (Number of spared neurons after secondary degeneration)/(Number of spared neurons after primary damage) x 100.

#### 7.1.2 ELECTROPHYSIOLOGICAL RECORDINGS

Nerves were excised and their compound action potentials (CAPs) were recorded *in vitro* using a suction electrode experimental set-up (Yoles et al., J. Neurotrauma 13:49-57, 1996). At different times after injury and injection of T cells or PBS, rats were killed by intraperitoneal injection of pentobarbitone (170 mg/kg) (CTS Chemical Industries, Israel). Both optic nerves were removed while still attached to the optic chiasma, and were immediately transferred to a vial containing a fresh salt solution consisting of 126 mM NaCl, 3 mM KCl, 1.25 mM  $\text{NaH}_2\text{PO}_4$ , 26 mM  $\text{NaHCO}_3$ , 2 mM  $\text{MgSO}_4$ , 2 mM  $\text{CaCl}_2$ , and 10 mM D-glucose, aerated with 95%  $\text{O}_2$  and 5%  $\text{CO}_2$  at room temperature. After 1 hour, electrophysiological recordings were made. In the injured nerve, recordings were made in a segment distal to the injury site. This segment contains axons of viable retinal ganglion cells that have escaped both primary and secondary damage, as well as the distal stumps of non-viable retinal ganglion cells

that have not yet undergone Wallerian degeneration. The nerve ends were connected to two suction Ag-AgCl electrodes immersed in the bathing solution at 37°C. A stimulating pulse was applied through the electrode, and the CAP was recorded by the distal electrode. A stimulator (SD9; Grass Medical Instruments, Quincy, Massachusetts) was used for supramaximal electrical stimulation at a rate of 1 pps to ensure stimulation of all propagating axons in the nerve. The measured signal was transmitted to a microelectrode AC amplifier (model 1800; A-M Systems, Everett, Washington). The data were processed using the LabView 2.1.1 data acquisition and management system (National Instruments, Austin, Texas). For each nerve, the difference between the peak amplitude and the mean plateau of eight CAPs was computed and was considered as proportional to the number of propagating axons in the optic nerve. The experiments were done by experimentors "blinded", to sample identity. In each experiment the data were normalized relative to the mean CAP of the uninjured nerves from PBS-injected rats,

### 7.1.3 CLINICAL EVALUATION OF EXPERIMENTAL AUTOIMMUNE ENCEPHALOMYELITIS

Clinical disease was scored every 1 to 2 days according to the following neurological scale: 0, no abnormality; 1, tail atony; 2, hind limb paralysis; 3, paralysis extending to thoracic spine; 4, front limb paralysis; 5, moribund state.

## 7.2 RESULTS

### 7.2.1 NEUROPROTECTION BY AUTOIMMUNE anti-MBP T CELLS

Morphological analyses were done to assess the effect of the T cells on the response of the nerve to injury, and specifically on secondary degeneration. Rats were injected intraperitoneally immediately after optic nerve injury with PBS or with  $1 \times 10^7$  activated T cells of the various cell lines. The degree of primary damage to the optic nerve axons and their attached RGCs was measured by injecting the dye 4-Di-10-Asp

distal to the site of the lesion immediately after the injury. A time lapse of 2 weeks between a moderate crush injury and dye application is optimal for demonstrating the number of still viable labeled neurons as a measure of secondary degeneration, and as the response of secondary degeneration to treatment. Therefore, secondary degeneration was quantified by injecting the dye immediately or 2 weeks after the primary injury, and calculating the additional loss of RGCs between the first and the second injections of the dye. The percentage of RGCs that had survived secondary degeneration was then calculated. The percentage of labeled RGCs (reflecting still-viable neurons) was significantly greater in the retinas of the rats injected with anti-MBP T cells than in the retinas of the PBS-injected control rats (Fig. 2). In contrast, the percentage of labeled 30 RGCs in the retinas of the rats injected with anti-OVA or anti-p277 T cells was not significantly greater than that in the control retinas. Thus, although the three T cell lines accumulated at the site of injury, only the MBP-specific autoimmune T cells had a substantial effect in limiting the extend of secondary degeneration. Labeled RGCs of injured optic nerves of rats injected with PBS (Fig. 3A), with anti-p277 T cells (Fig. 3B) or with anti-MBP T cells (FIG. 3C) were compared morphologically using micrographs.

#### 7.2.2 CLINICAL SEVERITY OF EAE

Animals were injected i.p. with  $10^7$   $T_{MBP}$  cells with or without concurrent optic nerve crush injury. The clinical course of the rats injected with the  $T_{MBP}$  cells was evaluated according to the neurological paralysis scale. Each group contained 5-9 rats. The functional autoimmunity of the injected anti-MBP T cells was demonstrated by the development of transient EAE in the recipients of these cells. As can be seen in Fig. 4A, the course and severity of the EAE was not affected by the presence of the optic nerve crush injury.

#### 7.2.3 SURVIVAL OF RGCs IN NON-INJURED NERVES

Animals were injected i.p. with  $10^7$   $T_{MBP}$  cells or PBS. Two weeks later, 4-Di-10-Asp was applied to the optic nerves. After five days the retinal were excised and flat

mounted. Labeled RGCs from five fields (located at approximately the same distance from the optic disk), in each retina were counted and their average number per area ( $\text{mm}^2$ ) was calculated.

As can be seen in Fig. 4B, there is no difference in the number of surviving RGCs per area ( $\text{mm}^2$ ) in non-injured optic nerves of rats injected with anti-MBP T cells compared to in rats injected with PBS.

#### 7.2.4. NEUROPROTECTION BY T CELLS REACTIVE TO A CRYPTIC EPITOPE

To determine whether the neuroprotective effect of the anti-MBP T cells is correlated with their virulence, the effect of T cells reactive to a "cryptic" epitope of MBP, the peptide 51-70 (p51-70) was examined. "Cryptic" epitopes activate specific T cells after an animal is immunized with the particular peptide, but not with the whole antigen (Mor et al., J. Immunol. 155:3693-3699. 1995). The T cell line reactive to the whole MBP and the T cell line reactive to the cryptic epitope p51-70 were compared for the severity of the EAE they induced, and for their effects on secondary degeneration. In rats injected with the T cell line reactive to the cryptic epitope, disease severity (as manifested by the maximal EAE score) was significantly lower than that in rats injected with the T cell line reactive to the whole protein (Table 1). Whereas anti-MBP T cells caused clinical paralysis of the limbs, rats injected with the anti-p51-70 T cells developed only tail atony, not hind limb paralysis, and almost none showed weakness of the hind limbs. Despite this difference in EAE severity, the neuroprotective effect of the less virulent (anti-p51-70) T cells was similar to that of the more virulent (anti-MBP) T cells (Fig. 5). The percentage of RGCs surviving secondary degeneration in the retinas of rats injected with either of the lines was significantly higher than in the retinas of the PBS-injected rats. Thus, there was no correlation between the neuroprotective effect of the autoimmune T cells and their virulence. It is possible that the anti-p51-70 T cells encounter little antigen in the intact CNS, and therefore cause only mild EAE. Their target antigen

may however become more available after injury, enabling these T cells to exert a neuroprotective effect.

TABLE 1. Anti-MBP and anti-p51-70 T cells  
Vary in Pathogenicity

<u>T Cell Line</u>	<u>Clinical EAE</u>	<u>Mean Max. Score</u>
Whole MBP	Moderate to severe	2.00 + 0.2
p51-70 of MBP	Mild	0.70 + 0.2

Immediately after optic nerve crush injury, Lewis rats were injected with activated anti-MBP T cells or anti-p51-70 T cells. The clinical course of EAE was evaluated according to the neurological paralysis scale. The mean maximal (max.) score  $\pm$  s.e.m. was calculated as the average maximal score of all the diseased rats in each group. The table is a summary of nine experiments. Each group contains five to ten rats. Statistical analysis showed a significant difference between the mean maximal score of rats injected with anti-MBP T cells and that of rats injected with anti-p51-70 T cells ( $P=0.039$ , Student's t-test).

#### 7.2.5 ELECTROPHYSIOLOGICAL ACTIVITY

To confirm the neuroprotective effect of the anti-MBP T cells, electrophysiological studies were done. Immediately after optic nerve injury, the rats were injected intraperitoneally with PBS or with  $1 \times 10^7$  activated anti-MBP or anti-OVA T cells. The optic nerves were excised 7, 11 or 14 days later and the compound action potentials (CAPs), a measure of nerve conduction, were recorded from the injured nerves. On day 14, the mean CAP amplitudes of the distal segments recorded from the injured nerves obtained from the PBS-injected control rats were 33% to 50% of those recorded from the rats injected with the anti-MBP T cells (Fig. 6A, Table 2). As the distal segment of the injured nerve contains both neurons that escaped the primary insult and injured neurons that have not yet degenerated, the observed neuroprotective effect could reflect the rescue of spared neurons, or a delay of Wallerian degeneration of the injured neurons (which normally occurs in the distal stump), or both. No effect of the injection anti-MBP T cells on the mean CAP amplitudes of uninjured nerves was



observed (Fig. 6B, Table 2). It is unlikely that the neuroprotective effect observed on day 14 could have been due to the regrowth of nerve fibers, as the time period was too short for this.

The strong neuroprotective effect of the anti-MBP T cells seen on day 14 was associated with a significantly decreased CAP amplitude recorded on day 7 (Table 2). The anti-MBP T cells manifested no substantial effect on the uninjured nerve on day 7, indicating that the reduction in electrophysiological activity observed in the injured nerve on day 7 might reflect the larger number of T cells present at the injury site relative to the uninjured nerve (Fig. 1). The observed reduction in CAP amplitude in the injured nerve on day 7 reflected a transient resting state in the injured nerve. This transient effect has not only disappeared, but was even reversed by day 14 (Table 2). Early signs of the neuroprotective effect could already be detected on day 11 in the rats injected with anti-OVA T cells, no reduction in CAP amplitude on day 7 could be detected in either the injured or the uninjured nerves, and no neuroprotective effect was observed on day 14 (Table 2). Thus, it seems that the early reduction in CAP and the late neuroprotection shown specifically by the anti-MBP T cells are related.

TABLE 2. Transient reduction in electrophysiological activity of the injured optic nerve induced by anti-MBP T cells, followed by a neuroprotective effect

	<u>Uninjured Optic Nerve</u>		<u>Injured Optic Nerve</u>	
	<u>Day 7</u>	<u>Day 14</u>	<u>Day 7</u>	<u>Day 14</u>
Ratio (%) $T_{\text{MBP}}/\text{PBS}$	89.9±9.4 (n=22)	101.2±22.7 (n=10)	63.8*±14.9 (n=17)	243.1**±70.8 (n=8)
Ratio (%) $T_{\text{OVA}}/\text{PBS}$	109.7±13.2 (n=11)	92.5±12.6 (n=3)	125.5±24.4 (n=11)	107.3±38.9 (n=4)

Immediately after optic nerve injury, rats were injected with PBS or with activated anti-MBP or anti-OVA T cells. After 7 or 14 days, the CAPs of injured and uninjured nerves were recorded. Ratios were calculated for uninjured nerves as (mean CAP of uninjured nerves from T cell-injected rats/mean CAP of uninjured nerves from PBS-injected rats) x 100, or for injured

nerves as (mean CAP of injured nerves from T cell-injected rats/mean CAP of injured nerves from PBS-injected rats) x 100. The P value was calculated by comparing the logarithms of the normalized CAP amplitudes of nerves from PBS-injected rats and rats injected with T cells, using the unpaired Student's test, \*P<0.05; \*\*P<0.001 n=sample size.

---

### 7.3 NEUROPROTECTION IN SPINAL CORD INJURY

#### 7.3.1. MATERIALS AND METHODS

Animals, antigens (MBP, OVA) and T cell lines were as described hereinbefore in 6.1.1, 6.1.3 and 6.1.4, respectively

**Contusion.** Adult rats (300 to 350g) were anesthetized and the spinal cord was exposed by laminectomy at the level of T7-T8. One hour after induction of anesthesia, a 10 gram rod was dropped onto the laminectomized cord from a height of 50 mm. The impactor device (designed by Prof. Wise Young) allowed, for each animal, measurement of the trajectory of the rod and its contact with the spinal cord to allow uniform lesion. Within an hour of the contusion, rats were injected i.p., on a random basis, with either 10<sup>7</sup> cells (specific to either MBP or OVA, depending on the experimental design) or with PBS. Bladder expression was done at least twice a day (particularly during the first 48h after injury, when it was done 3 times a day) until the end of the second week, by which time the rats had developed autonomous bladder voidance. Approximately twice a week, locomotor activity (of the trunk, tail and hind limbs) in an open field was evaluated by placing the rat for 4 min in the middle of a circular enclosure made of molded plastic with a smooth, non-slip floor (90 cm diameter, 7 cm wall height).

#### 7.3.2 RESULTS

The present study of spinal cord neuroprotection was prompted by the previous example that partial injury to an optic nerve can be ameliorated administering T cells directed to a CNS self-antigen. The question was whether autoimmune T cells could have a beneficial effect on recovery from traumatic spinal cord injury with its greater mass of injured CNS tissue and the attendant spinal shock.

Adult Lewis rats were subjected to a calibrated spinal cord contusion produced by dropping a 10 gram weight from a height of 50 mm onto the laminectomized cord at the level of T7-T8 (see description included in Basso et al., Exp-Neurol 139, 244-256, 1996). The rats were then injected intraperitoneally with autoimmune T cells specific to MBP. Control rats were similarly injured but received either no T cells or T cells specific to the non-self antigen ovalbumin (OVA). Recovery of the rats was assessed every 3 to 4 days in terms of their behavior in an open-field locomotion test, in which scores range from 0 (complete paraplegia) to 21 (normal mobility). The locomotor performance of the rats was judged by observers blinded to the identity of the treatment received by the rats. Included in the study was a group of uninjured, sham-operated (laminectomized but not contused) rats which were injected with anti-MBP T cells to verify the activity of the T cells. In all the sham-operated rats, the anti-MBP T cells induced clinical experimental autoimmune encephalomyelitis (EAE), which developed by day 4, reached a peak at day 7 and resolved spontaneously by day 11. Note, therefore, that at the early post-traumatic stage, any effect of the autoimmune T cells on the injured spinal cord, whether positive or negative, would be transiently masked both by spinal shock and by the paralysis of EAE.

Indeed, none of the rats with contused spinal cords showed any locomotor activity in the first few days after the contusion (Fig. 7A). Interestingly, however, the rats treated with anti-MBP T cells recovered earlier from spinal shock; on day 11, for example, when no recovery could be detected in any of the untreated control rats, significant improvement was noted in the T cell-treated rats (Fig. 7A). At all time points thereafter, the rats that had received the autoimmune T cells showed better locomotor recovery than did the untreated injured rats (Fig. 7A). Thus the autoimmune T cells, in spite of being encephalitogenic, did confer significant neuroprotection. Moreover, the phase of neuroprotective activity coincided with the phase of immune paralysis, supporting our suggestion that neuroprotection might be related to transient paralysis.

By one month after trauma the rats in both groups had reached a maximal behavioral score, which then remained at plateau for at least 3 months of follow-up. In the untreated rats, maximal recovery of locomotor behavior, as noted in previous reports of similarly severe contusion (Basso et al., *supra*), was marked by some ineffectual movement of hind-limb joints, but the rats showed no ability support their body weight and walk, and obtained a score of  $7.3 \pm 0.8$  (mean  $\pm$  SEM). In contrast, the average score of the rats that had been treated with the anti-MBP T cells was  $10.2 \pm 0.8$ , and in some rats the value was high as 13. All the rats in the treated group could support their body weight and some could frequently walk in a coordinated fashion. The difference between the two groups, based on 2-factor repeated ANOVA, was statistically significant ( $p < 0.05$ ). The recovery curve based on locomotor activity is nonlinear. The above-described increase in motor activity seen after treatment with the anti-MBP T cells could result from much higher percentage of spared tissue based on a linear regression curve on which the behavioral score is correlated with the amount of neural spinal cord tissue (for example, a difference between 11 and 7) on the locomotion score would be read as a difference between 30% and less than 10% of spared tissue).

In another set of experiments the rats were subjected to a more severe insult, resulting in a functional score of  $1.9 \pm 0.8$  (mean  $\pm$  SEM) in the untreated group and  $7.7 \pm 1.4$  in the treated group (Fig. 7B). This difference of more than 3 fold in behavioral scores was manifested by the almost total lack of motor activity in the control rats as compared with the ability of the autoimmune T cell-treated rats to move all their joints. The beneficial effect was specific to treatment with anti-MBP T cells; no effect was observed after treatment with T cells specific to the non-self antigen OVA (data not shown). The positive effect of the autoimmune T cells seems to be expressed in the preservation of CNS tissue that escaped the initial lesion, i.e., in neuroprotection. Therefore, the magnitude of the effect would be inherently limited by the severity of the insult; the more severe the lesion, the less the amount of spared tissue amenable to neuroprotection.

To determine whether clinical recovery could be explained in terms of preservation of spinal axons, we performed retrograde labeling of the descending spinal tracts by applying the dye rhodamine dextran amine (Brandt et al, J-Neurosci-Methods 45:35-40, 1992) at T12, below the site of damage. The number of dye-stained cells that could be counted in the red nucleus of the brain constituted a quantitative measure of the number of intact axons traversing the area of contusion. Sections of red nuclei from injured rats treated with anti-MBP T cells (Fig. 8) contained 5-fold more labeled cells than sections taken from the untreated injured rats. Photomicrographs of red nuclei taken from rats treated with anti-MBP T cells (with an open field score of 10) and from PBS-treated rats (with a score of 6) are shown in Fig. 8. These findings indicate that the reduction in injury-induced functional deficit observed in the T cell-treated rats can be attributed to the sparing of spinal tracts, resulting in a higher degree of neuron viability.

After a follow-up of more than 3 months, when the locomotor activity scores had reached a plateau, the site of injury of three of PBS-treated animals and three animals treated with anti-MBP T cells were analyzed by diffusion-weighted MRI. The cords were excised in one piece from top to bottom and were immediately placed in fixative (4% paraformaldehyde). Axial sections along the excised contused cord were analyzed. Fig. 9 shows the diffusion anisotropy in axial sections along the contused cord of a rat treated with autoimmune T cells, as compared with that of PBS-treated control rat. The images show anisotropy in the white matter surrounding the grey matter in the center of the cord. Sections taken from the lesion sites of PBS-treated control rats show limited areas of anisotropy, which were significantly smaller than those seen at comparable sites in the cords of the rats treated with the anti-MBP T cells. Quantitative analysis of the anisotropy, reflecting the number of spared fibers, is shown in Fig. 9. The imaging results show unequivocally that, as a result of the treatment with the autoimmune anti-MBP T cells, some spinal cord tracts had escaped the degeneration that would otherwise have occurred.

### 7.3.3 DISCUSSION OF RESULTS

No cure has yet been found for spinal cord lesions, one of the most common yet devastating traumatic injuries in industrial societies. It has been known for more than 40 years that CNS neurons, unlike neurons of the peripheral nervous system, possess only a limited ability to regenerate after injury. During the last two decades, attempts to promote regeneration have yielded approaches that lead to partial recovery. In the last few years it has become apparent that, although most of the traumatic injuries sustained by the human spinal cord are partial, the resulting functional loss is nevertheless far worse than could be accounted for by the severity of the initial insult; the self-propagating process of secondary degeneration appears to be decisive.

A substantial research effort has recently been directed to arresting injury-induced secondary degeneration. All attempts up to now have been pharmacologically based, and some have resulted in improved recovery from spinal shock. The present study, in contrast, describes a cell therapy that augments what seems to be a natural mechanism of self-maintenance and leads, after a single treatment, to long-lasting recovery. The extent of this recovery appears to exceed that reported using pharmacological methods.

In most tissues, injury-induced damage triggers a cellular immune response that acts to protect the tissue and preserve its homeostasis. This response has been attributed to macrophages and other cells comprising the innate arm of the immune system. Lymphocytes, which are responsible for adaptive immunity, have not been thought to participate in tissue maintenance. Adaptive immunity, according to traditional teaching, is directed against foreign dangers. Our studies now show, however, that the adaptive T cell immune response can be protective even when there is no invasion by foreign pathogens. In the case of tissue maintenance, the specificity of the T cells is to tissue self-antigens.

Our observation of post-traumatic CNS maintenance by autoimmune T cells suggests that we might do well to reevaluate some basic concepts of autoimmunity. T cells that are specific

to CNS self antigens in general, and to MBP in particular, have long been considered to be only detrimental to health. In the present study, however, the same T cell preparation that can produce EAE in the undamaged CNS was found to be neuroprotective in the damaged spinal cord, suggesting that the context of the tissue plays an important part in determining the outcome of its interaction with T cells. It would seem that the tissue deploys specific signals to elicit particular T cell behaviors. Among such signals are costimulatory molecules, particularly members of the B7 family (Lenschow et al., Annu. Rev. Immunol. 14:233-258, 1996). As shown hereinafter, the injured rat optic nerve transiently expresses elevated levels of the costimulatory molecule B7.2, which is constitutively expressed at low levels in the rat CNS white matter and which is thought to be associated with regulation of the cytokine profile of the responding T cells (H. L. Weiner, Annu. Rev. Med. 48:341-51, 1997). The early post-injury availability of the exogenous anti-MBP T cells, coinciding with the observed early post-injury increase in B7.2 would support the idea that signals expressed by the tissue might modulate the T cell response. It is thus conceivable that anti-MBP T cells which cause a monophasic autoimmune disease upon interacting with a healthy CNS nerve, might implement a maintenance program when they interact with damaged CNS tissue expressing increased amounts of B7.2 and probably other costimulatory molecules. The neuroprotective effects of the T cells may be mediated, at least in part, by antigen-dependent regulation of specific cytokines or neurotrophic factors (M. Kerschensteiner et al., J. Exp. Med. 189:865-870, 1999) produced locally at the site of injury.

Thus, the present invention is also directed to manipulating B7.2 co-stimulatory molecule to prevent or inhibit neuronal degeneration and ameliorate the effects of injury to or disease of the nervous system. B7.2 molecule can be up-regulated for this purpose, using drugs or by genetic manipulation, without undue experimentation.

In a recent study, it was reported that injury to the spinal cord triggers a transient autoimmune response to MBP (Popovich et al., J. Neurosci. Res. 45:349-63, 1996). However,

whether that response is detrimental or beneficial remained an open question (Popovich et al, J. Comp. Neurol. 377:443-464, 1997). From our present data, it would appear that the activation of anti-MBP T cells could indeed be beneficial. However, a supplement of exogenous autoimmune T cells may be required to overcome the restrictions on immune reactivity imposed by the immune-privilege of the CNS (J. W. Streilein, Science 270:1158-1159, 1995). The finding that autoimmune response can be advantageous suggests that natural autoimmune T cells may have undergone positive selection during ontogeny, as proposed by the theory of the immunological homunculus (I. R. Cohen, Immunol. Today 13, 490-494 (1992), and are not merely a default resulting from the escape from negative selection of T cells that recognize self antigens (C. A. Janeway, Jr., Immunol. Today 13:11-6, 1992). Such a response could then be considered as a mechanism of potential physiological CNS self-maintenance, which is, however, not sufficient for the purpose because of the immune-privileged character of the CNS.

A single injection of autoimmune T cells lasted for at least 100 days. Thus, this procedure offers a form of self-maintenance. This specific autoimmune response, when properly controlled, is useful as part of a self-derived remedy for spinal cord injury.

#### EXAMPLE: NEUROPROTECTIVE EFFECTS OF NS-SPECIFIC ANTIGEN

##### 8.1 MATERIALS AND METHODS

Animals, crush injury of rat optic nerve, and retrograde labeling are described above in Sections 6 and 7. A peptide based on amino acids 35-55 of myelin/oligodendrocyte glycoprotein (MOG p35-55) was chemically synthesized at the Weizmann Institute, Israel.

##### 8.1.1 INHIBITION OF SECONDARY DEGENERATION

Rats were injected intradermally in the footpads with MOG p35-55 (50 µg/animal) and IFA, or PBS ten days prior to optic nerve crush injury. Retinal ganglion cells were assessed two weeks after injury using retrograde labeling as described above. The number of RGCs in rats injected with PBS or MOG



p35-55 was expressed as a percentage of the total number of neurons in rats injected with MOG p35-55 in the absence of crush injury.

## 8.2 RESULTS

As shown in Fig. 10, the number of labeled retinal ganglion cells (indicating viable axons) was about 12.5 fold greater in animals injected with MOG p35-55 compared to animals receiving PBS.

### EXAMPLE: NEUROPROTECTIVE EFFECTS OF MBP ADMINISTERED ORALLY

## 9.1 MATERIALS AND METHODS

Animals, crush injury of rat optic nerve, and retrograde labeling of RGCs are described above in Sections 6 and 7.

### 9.1.1 INHIBITION OF SECONDARY DEGENERATION

Bovine MBP (Sigma, Israel) (1 mg/dose) was administered to rats by gavage using a blunt needle. MBP was administered 5 times, every third day, beginning 2 weeks prior to optic nerve crush injury. The number of RGCs in treated animals was expressed as a percentage of the total number of neurons in animals subjected to optic nerve crush injury but which did not receive MBP.

## 9.2 RESULTS

As shown in Fig. 11, the number of labeled RGCs was about 1.3 fold greater in animals treated with MBP compared to untreated animals.

### 9.3 THE B7.2 COSTIMULATORY MOLECULE IS ASSOCIATED WITH POST-TRAUMATIC MAINTENANCE OF THE OPTIC NERVE BY ORAL ADMINISTRATION OF MBP

#### 9.3.1 INTRODUCTION

Autoimmune T cells can under under certain conditions be beneficial to traumatized CNS axons. The effect of such T cells on the damaged tissue might be influenced by the nature

and amount of the costimulatory molecules it expresses. We show that the B7.2 costimulatory molecule is constitutively expressed in the intact rat optic nerve, and after injury is up-regulated at the margins of the injury site. Pre-injury induction of oral tolerance to MBP resulted in a further post-injury increase in B7.2 at the margins and at the injury site itself, as well as a better preservation of the traumatized nerve. Thus, B7.2 expression in the brain and its up-regulated after trauma seem to be directly related to post-traumatic maintenance displayed by autoimmune T cells.

Neuronal injury in the CNS causes degeneration of directly damaged fibers as well as of fibers that escaped the primary insult. It also triggers a systemic response of autoimmune T cells to MBP, that might affect the course of degeneration of the injured nerve. Whether the effect of these T cells on the nerve is detrimental or beneficial may depend, in part, on the nature and level of the costimulatory molecules expressed by the damaged tissue. Several costimulatory molecules have recently been identified, including the B7 and CD40 molecules (Caux et al., "Activation of Human Dendritic Cells Through CD40 Cross-Linking", J. Exp. Med. 180:1263-1272, 1994; and Lenschow et al., "CD28/B7 System of T Cell Costimulation", Annu. Rev. Immunol. 14:233-258, 1996). CD40 appears to be dominant during cell differentiation in the lymph nodes and B7 during activation of T cells in the target organ (Grewal et al., "Requirement for CD40 Ligand in Costimulation Induction, T Cell Activation, and Experimental Allergic Encephalomyelitis", Science 273:1864-1867, 1996). B7 costimulatory molecules are expressed on antigen-presenting cells (APCs) as B7.1 or B7.2., which might preferentially support activation of the Th1 or the Th2 type of immune response, respectively (Kuchroo et al., "B7-1 and B7-2 costimulatory molecules activate differentially the Th1/Th2 developmental pathways: application to autoimmune disease therapy", Cell 80:707-718, 1995; and Karandikar et al., "Targeting the B7/CD28:CTLA-4 costimulatory system in CNS autoimmune disease", J. Neuroimmunol. 89:10-18, 1998). We were therefor interested in determining the identity B7 subtype expressed in intact and injured CNS white matter, and its

possible influence on the course of the response to the injury.

### 9.3.2 RESULTS

The costimulatory molecule expressed constitutively in the intact optic nerves of adult Lewis rats was identified as B7.2. (Figs. 12A, 12B). To examine the effects of neurotrauma on the expression of B7 costimulatory molecules, we inflicted a mild crush injury on the optic nerves of Lewis rats and assessed the neural expression of B7 by immunohistochemical analysis. The most striking effect of the injury was seen on B7.2 expression manifested on post-injury day 3 by its elevation at the margins of the injury site (Figs. 12C,D,E). In contrast, expression of B7.1 was not detected in the optic nerve either before or 3 days after injury. On day 7, however, B7.1 was detectable at the site of injury, having pattern reminiscent of that seen for macrophages or microglia (Fig. 12F).

Next, we attempted to determine whether the degenerative response to optic nerve injury could be modified by peripheral manipulation of the immune system. The manipulation chosen was induction of oral tolerance, known to cause a "bystander" T cell immunosuppressive effect (Weiner et al., "Tolerance Immune Mechanisms and Treatment of Autoimmune Diseases", Immunol. Today 18:335-343, 1997). Ingestion of low doses of MBP results in the activation of T cells which, based on antigen recognition, secrete TGF as the dominant cytokine and thus favor an immune response of Th2/3 type (Chen, Y., "Regulatory T Cell Clones Induced by Oral Tolerance: Suppression of Autoimmune Encephalomyelitis", Science 265: 1237-1240, 1994).

Lewis rats were fed with food to which 1 mg of bovine MBP had been added five times daily every other day. Ten days after first receiving the supplement, the rats were subjected to mild unilateral optic nerve crush injury. This time interval between initiation of oral tolerance and injury was chosen to allow adequate build-up of the systemic T cell response. As shown in Fig. 13A and B, the numbers of macrophages or active microglia (indicated by ED-1 labeling)

and T cells (indicated by immunolabeling for T cell receptor), assessed 3 days after injury, did not differ from those observed in control injured rats which did receive any treatment or were fed with PBS. In the rats with induced oral tolerance to MBP, however, the amounts B7.2 were further increased at the margins of the site of injury (Fig. 13C) as compared with controls (Fig. 12E). In addition, B7.2 in the rats with induced oral tolerance to MBP was also elevated at the site of injury relative to the control nerves (Fig. 13C). It seems reasonable to assume that the T cells exposed to MBP via intestinal absorption, upon invading the injured CNS, contributed to the increase in expression of B7.2 by the injured nerve.

We then attempted to determine whether the observed changes in B7.2 expression in the injured rats was correlated with the extent of neuronal degeneration. Acute injury of the rat optic nerve is followed by a process of nerve degeneration, which can be quantified by retrograde labeling of the surviving neurons and counting of the corresponding cell bodies. Two weeks after optic nerve injury the number of surviving retinal ganglion cells (RGCs), representing still-viable neurons, in the group of MBP-fed rats was significantly higher than that in the control group, or than in the group of rats with injured nerves that were fed with ovalbumin. Interestingly, the benefit of the induced oral tolerance to MBP was increased by feeding the rats with more intensive schedule (Fig. 14).

#### DISCUSSION OF EXPERIMENTAL RESULTS

The results of the experiments described in Sections 6 and 7 show that activated T cells accumulate at a site of injury in the CNS. Furthermore, the results also demonstrate that the accumulation of T cells at the site of injury is a non-specific process, i.e., T cells which accumulated at the site of injury included both T cells which are activated by exposure to an antigen present at the site of injury as well as T cells which are activated by an antigen not normally present in the individual.

The results of experiments described in Section 7 demonstrate that the beneficial effects of T cells in

ameliorating damage due to injury in the CNS are associated with an NS-specific self-antigen as illustrated by MBP. More specifically, the administration of non-recombinant T cells which were activated by exposure to an antigen which can cause autoimmune disease ( $T_{MBP}$ ), rather than aggravating the injury, led to a significant degree of protection from secondary degeneration. Thus, activating T cells by exposure to a fragment of an NS-specific antigen was beneficial in limiting the spread of injury in the CNS. The present findings show that secondary degeneration can be inhibited by the transfer into the individual on non-recombinant T cells which recognize an NS-specific self antigen which is present at a site of injury. The T cells may recognize cryptic or non-pathogenic epitopes of NS-self antigens.

In addition, the studies described in Sections and 9 show that activation of T cells by administering an immunogenic antigen (e.g. MBP) or immunogenic epitope of an antigen (e.g. MOG p35-55), may be used for preventing or inhibiting secondary CNS degeneration following injury.

The foregoing description of the specific embodiments will so fully reveal the general nature of the invention that others can, by applying current knowledge, readily modify and/or adapt for various applications such specific embodiments without undue experimentation and without departing from the generic concept, and, therefore, such adaptations and modifications should and are intended to be comprehended within the meaning and range of equivalents of the disclosed embodiments. It is to be understood that the phraseology or terminology employed herein is for the purpose of description and not of limitation. The means, materials, and steps for carrying out various disclosed functions may take a variety of alternative forms without departing from the invention. Thus the expressions "means to..." and "means for...", or any method step language, as may be found in the specification above and/or in the claims below, followed by a functional statement, are intended to define and cover whatever structural, physical, chemical or electrical element or structure, or whatever method step, which may now or in the future exist which carries out the recited function, whether or not precisely equivalent to

the embodiment or embodiments disclosed in the specification above, i.e., other means or steps for carrying out the same function can be used; and it is intended that such expressions be given their broadest interpretation.

All publications cited herein are incorporated by reference in their entirety.

WHAT IS CLAIMED IS:

1. A composition for preventing or inhibiting degeneration in the central nervous system or peripheral nervous system for ameliorating the effects injury or disease, comprising:
  - (a) NS-specific activated T cells;
  - (b) NS-specific antigen;
  - (c) a peptide derived from an NS-specific antigen;
  - (d) a nucleotide sequence encoding an NS-specific antigen;
  - (e) a nucleotide sequence encoding a peptide derived from an NS-specific antigen, or
  - (f) any combination of (a)-(e).
2. A composition according to claim 1, for promoting nerve regeneration in the central nervous system or peripheral nervous system for ameliorating the effects of injury or disease.
3. The composition of claim 1 or 2 in which said injury comprises spinal cord injury, blunt trauma, penetrating trauma, hemorrhagic stroke, or ischemic stroke.
4. The composition of claim 1 or 2 in which said disease is Diabetic neuropathy, senile dementia, Alzheimer's disease, Parkinson's Disease, facial nerve (Bell's) palsy, glaucoma, Huntington's chorea, amyotrophic lateral sclerosis, non-arteritic optic neuropathy, or vitamin deficiency.
5. The composition of claim 1 or 2 in which said disease is not an autoimmune disease or a neoplasm.
6. The composition according to any of of claims 1-5 wherein said NS-specific activate T cells of (a) are autologous T cells, or allogeneic T cells from related donors, OR HLA-matched or partially matched, semi-allogeneic or fully allogeneic donors.
7. The composition according to claim 6 wherein said autologous T cells have been stored or are derived from autologous CNS cells.
8. The composition according to claim 6 wherein said T cells are semi-allogeneic T cells.

9. The composition according to any of claims 1-5 wherein said NS-specific antigen of (b) is elected from myelin basic protein (MBP), myelin oligodendrocyte glycoprotein (MOG), proteolipid protein (PLP), myelin-associated glycoprotein (MAG), S-100,  $\beta$ -amyloid, Thy-1, PC, P2 and neurotransmitter receptors.

10. The composition according to any one of claims 1-5 wherein said peptide derived from an NS-specific antigen is an immunogenic epitope or a cryptic epitope of said antigen.

11. The composition according to claim 10 wherein said peptide is an immunogenic epitope or a cryptic epitope derived from MBP.

12. The composition according to claim 11 wherein said peptide corresponds to the sequences p11, p51-70, p91-110, p131-150, or p151-170 of MBP.

13. The compositions according to any one of claims 1-5 and 11-12 in which said NS-specific antigen or a peptide derived therefrom is administered intravenously, orally, intranasally, intrathecally, intramuscularly, intradermally, topically, subcutaneously, mucosally (e.g., orally, intranasally, vaginally, rectally) or buccally.

14. The composition according to claim 13 comprising MBP for oral administration.

15. Use of:

- (a) NS-specific activated T cells;
- (b) an NS-specific antigen;
- (c) a peptide derived from an NS-specific antigen;
- (d) a nucleotide sequence encoding an NS-specific

antigen;

(e) a nucleotide sequence encoding a peptide derived from an NS-specific antigen, or

(f) any combination of (a)-(e),

for the preparation of a composition for preventing or inhibiting neuronal degeneration in the central nervous system or peripheral nervous system for ameliorating the effects of injury or disease.

16. A method for preventing or inhibiting neuronal degeneration in the central nervous system or peripheral



nervous system, which comprises administering to an individual in need thereof an effective amount of:

- (a) NS-specific activated T cells;
- (b) NS-specific antigen;
- (c) a peptide derived from an NS-specific antigen;
- (d) a nucleotide sequence encoding an NS-specific antigen;
- (e) a nucleotide sequence encoding a peptide derived from an NS-specific antigen, or
- (f) any combination of (a)-(e).

17. A method for preventing or inhibiting neuronal degeneration in the central nervous system or peripheral nervous system comprising administering to an individual in need thereof an effective amount of a composition according to any one of claims 1-13 and actively immunizing said individual to build up a critical T cell response.

18. A method for preventing or inhibiting neuronal degeneration in the central nervous system or peripheral nervous system comprising administering to an individual in need thereof an effective amount of a composition for up-regulating B7.2 costimulatory molecule or genetically manipulating B7.2 costimulatory molecule in said individual.




FIG. 1




FIG. 2




FIG. 3C



FIG. 3B




FIG. 3A




FIG. 4A




FIG. 4B




FIG. 5




FIG. 6






FIG. 7 A




FIG. 7 B

FIG. 8





FIG. 9






FIG. 10




FIG. 4

FIG. 12




FIG. 13





FIG. 14



Experiment Groups



```
1 ccaagaagat cccacagcag cttccgaagg cctggatgtg atggcatcac agaagagacc
61 ctcacagcga cacggatcca agtacttggc cacagcaagt accatggacc atgcccggca
121 tggttctctc ccaaggcaca gagacacggg catccttgac tccatcgggc gcttctttag
181 cggtgacagg ggtgcgcca agcggggctc tggcaaggac tcacacaca gaactacca
241 ctacggctcc ctgccccaga agtcgcagag gaccaagat gaaaaccag tagtccactt
301 cttcaagaac attgtgacac ctcgtaacc cctccatcc caaggaaagg ggagaggcct
361 gtccctcagc agatttagct ggggaggaag agacagcgc tctggatctc ccatggcaag
421 acgtgagag cctccctgct cagccttccc gaatcctgcc ctcggcttct taatataact
481 gccttaaacg ttttaattcta cttgcaccaa atagctagtt agagcagacc ctctcttaat
541 cccgtggggc tgtgaacgcg gcgggccagc ccacggcacc ctgactggct aaaactgttt
601 gtcccttttt at
```

FIG. 15

1 gaaaacagtg cagccacctc cgagagcctg gatgtgatgg cgtcacagaa gagaccctcc  
61 cagaggcacg gatccaaagta cctggccaca gcaagtacca tggaccatgc caggcatggc  
121 ttcctcccaa ggcacagaga caggggcac cttgactcca tcgggcgctt ctttggcggg  
181 gacaggggtg cgccaaagcg gggctctggc aaggactcac accaccgggc aagaactget  
241 cactatggct ccctgcccga gaagtcacac ggccggaccc aagatgaaaa ccccgtagtc  
301 cactttctca agaacattgt gacgcctcgc acaccacccc cgtcgcaagg aaaggggaga  
361 ggactgtccc tgagcagatt tagctggggg gccgaaggcc agagaccagg atttggtac  
421 ggaggcagag cgtccgacta taaatcggt cacaagggat tcaaggagat cgatgcccag  
481 ggcacgcttt ccaaaatttt taagctggga ggaagagata gtcgctctgg atcaccatg  
541 gctagacgct gaaaaccac ctgggtccgg aatcctgtcc tcagcttctt aatataactg  
601 ccttaaaact ttaatccac ttgcccctgt tacctaatta gagcagatga cccctccct  
661 aatgcctgcg gagttgtgca cgtagtggg tcaggccacg gcagcctacc ggcaatttcc  
721 ggccaacagt taaatgagaa catgaaaaca gaaaacgggt aaaactgtcc ctttctgtgt  
781 gaagatcacg ttccttcccc cgcaatgtgc ccccgagcgc acgtgggtct tcagggggcc  
841 aggtgcacag acgtccctcc acgttcaccc ctccaccctt ggactttctt ttcgcccgtg  
901 ctcggcacc cttgcgtttt gctggctact gccatggagg cacacagctg cagagacaga  
961 gaggacgtgg gcggcagaga ggactgttga catccaagct tcctttgttt ttttttctg  
1021 tctttctctc acctcctaaa gtagacttca tttttcctaa caggattaga cagtcaagga  
1081 gtggcttact acatgtggga gctttttgt atgtgacatg cgggctgggc agctgttaga  
1141 gtccaaacgtg gggcagcaaa gagagggggc cacctcccca ggccgtggct gcccacacac  
1201 cccaattagc tgaattcgcg tgtggcagag ggaggaaaag gaggcaaacg tgggctgggc  
1261 aatggcctca cataggaaac aggttcttcc tggagatttg gtgatggaga tgtcaagcag  
1321 gtggcctctg gacgtcaccc ttgcccctga tgggtggccc agagcagcct ctatgaacaa  
1381 cctcgtttcc aaaccacagc ccacagccgg agagtccagg aagacttgcg cactcagagc  
1441 agaagggtag gagtctctca gacagcctcg cagccgcgc agtcgcccac agacactggc  
1501 tgtgaccggg cgtgctggca gcggcagtgc acagtggcca gcactaacc tccctgagaa  
1561 gataaccggc tcattcactt cctcccagaa gacgcgtggg agcagtagg cacaggcgtg  
1621 cactgtctcc cgaattactc accgagacac acgggctgag cagacggccc ctgtgatgga  
1681 gacaaagagc tcttctgacc atatccttct taacaccgc tggcatctcc tttcgcct  
1741 cctccctaa cctactgacc caccttttga ttttagcgca cctgtgattg ataggcctc  
1801 caaagagtcc cagctggca tcaccctccc cgaggacgga gatgaggat agtcagcgtg  
1861 atgccaaaac gcgctcttct aatccaattc taattctgaa tgtttcgtgt gggcttaata  
1921 ccatgtctat taatatatag cctcgatgat gagagagtta caaagaacaa aactccagac  
1981 aaaaacctcc aaatttttca gcagaagcac tctgcgtcgc tgagctgagg tcggctctgc  
2041 gatccatacg tggccgcacc cacacagcac gtgctgtgac gatggctgaa cggaaagtgt  
2101 acactgttcc tgaatattga aataaaacaa taaactttt

FIG. 16

A.

```

1 taatatctag ggktttgact ctgaccctgt ttggggctct cacttcctgg cttctcagc
61 ttgtgctgca tatccacac caattagacc caaggatcag ttggaagttt ccaggacatc
121 ttcattttat ttccaccctc aatccacatt tccagatgtc tctgcagcaa agcgaaattc
181 caggcaagcc ttagggaaaa aaggaaaaac aaagaaaatg aaacaatttg cagtgaagg
241 cagaaagaga agatggagcc cttagagaag ggagtatccc tgagtagggt gggaaaagg
301 gaggagaagg ggaggaggag aggaggaggga aagcaggcct gtccctttaa gggggttggc
361 tgtcaatcag aaagcccttt tcattgcagg agaagaggac aaagatactc agagagaaaa
421 agtaaaagac cgaagaaggga ggctggagag accaggatcc ttccagctga acaaagtcag
481 ccacaaagca gactagccag ccggtacaa ttggagtcag agtcccaaag acatgggtaa
541 gtttcaaaaa ctttagcatt gaagattcaa gaggacacag g

```

B.

```

1 ctgctttcag agcctgtgac ttcttgtgtg cctctcctgt ttctcagcaa catggcatag
61 ggccctggat accaggctctg gggatctcag ggactcttag cactttaaga cacatgtgtt
121 cccaggccct ggtgtgttcc tctagtgcga gaaagatgtt tcatgctttg ctgactttgt
181 ataaagtctg tttgtagctg ttttgacaga atctcagcgt ataactgagg gtggggacat
241 tagccaagct gcattatagg aggacaaaac tgccatacaa agtgtccaaa atcattaagc
301 ctgcattttt attattggga gtaatatcaa acctcctatt ttccaatttt catttcttgt
361 cctgtgctag ctccatcctg tttggactgc tcctcccata tgtaaaactaa gaagaatcaa
421 gcattctttg caacaaatc acacgatgtc caaaaatgtc caggagcctc caatttccaa
481 agtttctctc acctggaatg ctcttcacgc taaaatcctg tctgacaata ccagcatctc
541 tggcctgcac tcaccccttc ctggaactcc aagtgcattt acctctgttt accacttact
601 tggctgcctg aattgttagt tgaaaatatt aggtctactt agctaattct tctcaggaa
661 attaaagact cccatatggc agagtctgtg tcttttctct cttcatatcc cgtataacac
721 ccagcataat gctgggcata tagtgagtat tccataaata gttgatgaat gactaaaata
781 agcaagcaaa caaacagact agaacaataa gaaagaaggg actggatttc ataactctc
841 tggcttgcta tttgaattgc tgaattatta ttatttatta aatatttttt aaattctggc
901 aataaaaagg aaggatttat tttctttctt tctttttttt tttcttgaga cagagtctcg
961 ctcttactgc ccaggctgga gtacaatggc gcaatcttgg ctacaggcaa cctccgctc
1021 ctccctgggt taacagattc tcctgtctca gctcctgag tagctgggat tacaggcata
1081 cggccatgcc cggctaattt ttgtattttt agtagagacg gggttttgcc atgttgcca
1141 ggctggctt gaactcctga cctcatgtga tccacctgcc tcagcctccc aaagtgtcg
1201 gattacaggc atgcccacc gtgcccggcc aaagatttat ttcaagaat gaaacaaagt
1261 aaggattctg ggtcaatctc acatgctgaa agccaaaacc tctagccgct cctgctttt
1321 gacttcggag tgcccactat ctccagcct gtgagcacag ggccctggcag aggggtttga
1381 gtggcatgag ctacctactg gatgtgctg actgtttccc cttcttcttc cccaggctg
1441 ttagagtgtc gtgcaagatg tctggtaggg gccccctttg ctccctgggt ggccactgga
1501 ttgtgtttct ttgggggtggc actgttctgt ggctgtggac atgaagccct cactggaca
1561 gaaaagctaa ttgagacctt ttctccaaa aactaccaag actatgagta tctcatcaat
1621 gtgtaagtac ctgcccctcc acacagaccc atcttttttt tccctctctc catcctggag
1681 atagagaact cttcagtacc ttagtaacta gcaggggact ggggtggagc cagaccggat
1741 tcccagctct tccctctgtg ca

```

FIGS. 7A-E

C. 1 ctagaaaatc cctagccttg ttaaggtget cgetctggtg tatacctcac ttatgtcggg  
 61 aaagaagcca ggtcttcaat taataagatt ccctggtctc gtttgtctac ctgtaaatgc  
 121 aggatccatg ccttccagta tgtcatctat ggaactgcct ctttctctct cctttatggg  
 181 gccctcctgc tggctgaggg cttctacacc accggcgag tcaggcagat ctttggcgac  
 241 tacaagacca ccactctgcg caagggcctg agcgcaacgg taacaggggg ccagaagggg  
 301 aggggttcca gaggccaaaca tcaagctcat tctttggagc ggggtgtgtca ttgtttggga  
 361 aaatggctag gacatcccga caagggtgat atcctcagga ttttgtggca ataacaaggg  
 421 gtgggggaaa attgggcgcg agtctgtggc ctctgcccc cccaaggctg ggtcctctct  
 481 aggggccttg catttgagtg aggaagcgat ggctgcagcc gaacgagaag gtcaggaaga  
 541 acgtggtgcc cagctggctt agcctcacct ttcaaagggt ccctaagcaa atttctctc  
 601 aaaacagaaa gcatgagttt tgtgggatgc tttgtacaat cagaccattt ctaagccatc  
 661 tgttggatc cctttgttcc ctctctagta ggtaccacaa gagtggatct aactggacaa  
 721 gagtctaaaa tgctgctcat gtgattgaga cttgggcacc tgagctraga gggaggatgg  
 781 ataataaaaa ttaaataata actccaaggt aaatttaciaa tgttctgg

D. 1 gactctctc attcttcccc taccatttcc cccacccctc cgttatactg gggccagtta  
 61 tctagtagat actgccaatt acccttggca gaggtgccct gctcactaat tttatattggg  
 121 ggagmgccct ggaacctggg ttaaatgtct ggcacacgcc acttccagga tctcccagtt  
 181 tgtgtttcta catctgcagg ctgatgtga tttctaacca acccatgtca atcattttag  
 241 ttgtgggca tcacctatgc cctgaccgtt gtgtggctcc tgggtgttgc ctgctctgct  
 301 gtgctgtgt acatttactt caacacctgg accacctgcc agtctattgc cttccccagc  
 361 aagacctctg ccagtatagg cagtctctgt gctgatgcc gaatgtatgg tgagttaggg  
 421 tacgggtgct ttggctctcc taccactat ggaagcacta tatatttggg tattttctta  
 481 gtgtaaggag ggtggtgatt atgagaaaaa tataagatga tgaatgattg ggtcttagtt  
 541 tattaatcct tccctactga aaccagagag gtttcttccc ccggaaggga acttgggaagt  
 601 ggtgggagtt ttcttggcca ttacatttgg cctactctag ttgactgctg ttcacaaccc  
 661 caaagcagca catttcaata acaaacacaa ggttdsacca ctgttcaata ccaccttctc  
 721 ttttttgtaa acctgtagaa aagaggatcc taattgttgg tagmatccaa mtttacagcc  
 781 aggataatta gagatggaag aagggtctct ggggaaagtc tccatgtggc cccgtaactc  
 841 cataaagctt accctgcttg ctttttgtgt cttacttagg tgttctccca tggaaatgctt  
 901 tccctggcaa ggtttgtggc tccaaccttc tgtccatctg caaaacagct gaggtgagtg  
 961 ggttatttgg gttattttac aaggagtag ctaataccat acaaattaca cccatggcct  
 1021 tcaattttta ggactgaaag tttccctttg ctggattttg aattagccga ttgcttctta  
 1081 caacatgttg gctaagtgtg cctgagccaa tgagcataga aggtaaaaa cctcttttct

E. 1 aattagcaca cagaaaggat atccaacaca tacaaagctg tnntcatgga ctacactgga  
 61 gcatattact gctgttgcaa gaaacatttc ttcttctct tttcattttc ctgcagttcc  
 121 aaatgacctt ccacctgttt attgctgcat ttgtgggggc tgcagctaca ctggtttccc  
 181 tgggtgagttg actttgaatg atcttggcaa gtaaataggc ctgagatagt tgtgggtaca  
 241 gctatttctga aaggcaagaa ggtagactgc ttccatcctt gaaatgctgg agggga

FIGS. 17C-E

F. 1 aattctatat actatcacta tggctccact ttggatactc tccagtggat ttagttactc  
61 atattggaat acctgggagg acctcctaac attattagaa ttgttatgat tataatacaa  
121 ygcattgtcc caggctctgc tgatagtgc acagtgcct gtgaatgtag tgtgctcatt  
181 gtgcagatta aaaacctaaag gcactgaagg gtgaagtgat ttatctgaag ttattttata  
241 aagcagtgat cagacaasct gagctcacag aactccctgg cccctactgc tgagggttcc  
301 atacagagtc aagtaatttc tcacctgtta aaacgaattg attcattaac caggggagag  
361 ctctactgca tgatgtggct gtgtgtctac agcaagcacc ctatgactct aagtcactcg  
421 gacatattga tgtggcaaaag cccaaatatt gttcacttcc ctgaggaaaa ctcagtgtta  
481 gatcaaacag aggtgtggaa taaatcttta tgatttgatt ctctgggctt gggccatgag  
541 acctatgatg cctcagagac atcggacttc cagtcaagtgt tatatggaga aagccaagcc  
601 tgggatgtac tgctttttgc agagcatggg tttttccctt atttagttat gattttattt  
661 ctacccttcc tcattcccaa agggatttga ggaggggagtg ctttcttttc tactctcatt  
721 cacattctct cttctgttcc ctacagctca ccttcattgat tgctgccact tacaactttg  
781 ccgtccctaa actcatgggc cgaggcacca agttctgata ccccgtagaa atcccccttt  
841 ctctaataagc gaggtcttaa ccacacagcc tacaatgtctg cgtctcccat cttaactctt  
901 tgcctttgcc accaactggc cctctcttta cttgatgagt gtaacaagaa aggagagtct  
961 tgcagtgtat aaggtctctc tttggactct cccctcttat gtacctctt tagtcatttt  
1021 gcttcatagc tggttccctgc tagaaatggg aaatgcctaa taatatgact tcccaactgc  
1081 aagtcacaaa ggaatggagg ctctaattga attttcaagc atctcctgag gatcagaaaag  
1141 taatttcttc tcaaagggtta cttccactga tggaaaacaaa gtggaaggaa agatgctcag  
1201 gtacagagaa ggaatgtctt tggctctctt gccatctata ggggccaaat atattctctt  
1261 tgggtgtacaa aatggaattc attctgcgtc tctctattac actgaagata gaagaaaaaa  
1321 gaatgtcaga aaaacaataa gagcgtttgc ccaaatctgc ctattgcagc tgggagaagg  
1381 gggtcaaaagc aaggatcttt caccacaga aagagagcac tgaccccgat ggcgatggac  
1441 tactgaagcc ctaactcagc caaccttact tacagcataa gggagcgtag aatctgtgta  
1501 gacgaagggg gcatctggcc ttacacctcg ttagggaaga gaaacagggt cttgtcagca  
1561 tcttctcact ccttctctct tgataacagc taccatgaca accctgtggt ttccaaggag  
1621 ctgagaatag aaggaaaacta gcttacatga gaacagactg gcctgaggag cagcagttgc  
1681 tgggtggctaa tgggtgtaacc tgagatggcc ctctggtaga cacaggatag ataactcttt  
1741 ggcatacatg tcttttttct tgtaatttag ttgtgtactc tggcctctgt catatcttca  
1801 caatggtgct catttcatgg ggtattatcc attcagtcac cgtagggtgat ttgaagggtct  
1861 tgatttgttt tagaatgatg cacatttcat gtattccagt ttgtttatta cttatttggg  
1921 gttgcatcag aaatgtctgg agaataattc tttgattatg actgtttttt aaactaggaa  
1981 aattggacat taagcatcac aaatgatatt aaaaattggc tagttgaatc tattgggatt  
2041 tctacaagt attctgcctt tgcagaaaca gatttgggtga atttgaatct caatttgagt  
2101 aatctgatcg tcttttctag ctaattggaa atgattttac tttagcaatgt tatcttgggtg  
2161 tgtaagagt taggtttaac ataaagggtta ttttctctctg atatagatca cataacagaa  
2221 tgcaccagtc atcagctatt cagttggtaa gcttccagtc atcagctatt cagttggtaa  
2281 gcttcccagg aaaaaggaca ggcagaaaga gtttgagacc tgaatagctc ccagatttca  
2341 gtcttttaatt gtttttggta actttgggtt aaaaaaaaaa aaagtctgat tgggttttaatt  
2401 tgaaggaaaag atttgtacta cagttctttt gttgtaaaga gttgtgttgt tcttttcccc  
2461 caaagtgggt tcagcaatat ttaaggagat gtaagagctt tacaaaaaga cacttgatac  
2521 ttgttttcaa accagtatac aagataagct tccaggctgc atagaaggag gagagggaag  
2581 atgttttcta agaaaccaat caagataaag gacagtgaag taatccgtac cttgtgtttt  
2641 gttttgattt aataacataa caaataacca acccttccct gaaaacctca catgcataca  
2701 tacacatata tacacacaca aagagagtta atcaactgaa agtggttccct catttctgat  
2761 atagaattgc aattttaaca cacataaagg ataaactttt agaaacttat cttacaaagt  
2821 gtatttttat aaattaaaga aaataaaatt aagaatgttc tcaatcaaac atcgtgtcct  
2881 ttgagtgaat tgttctattt gacttcacaa tagaaactta ataactgtac cttctcaaqa

FIG. 17F

```

1  atggaaatgt  tctgtatattg  tgttgtctga  tgagataacc  actaactgta  gtgctattga
61  gcatttgaaa  catggctagt  gtaatcaatg  aaccaaattt  ttaattttat  ttaattgtaa
121 ttaattttta  gtggccacat  gcagggagtg  actgctgcat  tggacagcac  ggctctaaat
181 tgagcctttt  ttccttattt  ggtgagggat  acttgccctt  agattgggaa  gtctattttt
241 ggaacctgct  accaatgctg  gtctcacact  tgcaattctc  agctgagcca  agaggtgaga
301 gaaagggtcat  ttccatttcc  aagatctcac  tctccctgt  gacactgagg  aaactggcaa
361 gtgatgtgaa  ggctggagag  cgtgtccctg  atgctggctc  tgtcccttct  gcctgtgttg
421 actgacatag  ttagttgctg  cccttgctgg  tctcccttcc  tccaaccttg  cctctctgag
481 cacacctgac  attcatctca  tgacttccct  aaaaacattc  tttgggaaca  agaaactaac
541 aaatcccaa  tgacctatca  catatacaaa  catacagggc  agagtttga  ttccggtgag
601 aagaaaggga  ggttagacat  taagaagaat  ggtctggtga  tgacagtgtg  gagataatag
661 aaacaggaaa  aagaaatcta  agttttcttt  ctttttttaa  gaaccaataa  taatttctct
721 cttttgacta  gtcagttagg  ctggggtgga  ttggagggaag  cttacatatt  ccatgaacaa
781 gcctcttctc  aaggtcctgt  aagtgatect  gccccactga  ttagccctta  gaagaccctt
841 caaagggttg  atctccagga  gggagtgggg  gaggaagcc  ctgtaccagg  cagcctctgc
901 tccattgctc  tgggggggtg  gggagacaa  accctggtca  tccctcagt  ctgtagccct
961 tttgtgtgag  tgcctggcaa  ggggtgacgt  gggctgtttc  tgcgggcaca  gctgcagcaa
1021 ttaccggagt  ggaggcagg  ccaggcagc  actgccctcc  aagatcttcc  cttgggcttt
1081 tcagcagtaa  ggggacatgc  accccaagg  cctccacttg  gcctgacctt  gctgcggggg
1141 ctctctgtcc  ccaggaacag  tagagatgg  aagcttatcg  agacctctc  tgcccagctg
1201 cctctgtctc  ttctctctc  tctctctct  ccaagtgtct  tccagctatg  caggtaagac
1261 atgtttttt  tctgtccctg  gggagacct  gaaaacagaa  aggttagttt  cctgggggtt
1321 agctccttca  aacatcctca  agttggtata  ttatctttct  aaaacataga  cctactgaca
1381 tgcttccct  cctcagaaac  ctccgtggg  tggttcttac  agccttcaag  atggagtcca
1441 gactctttt  ttttttggg  acagagtctc  cctctgttgc  tcaggctgga  gtgcagtggc
1501 atgatctcgg  ctactgcaa  cctcagcctc  cctggttcaa  gcgattctcc  tgacttggcc
1561 tcccaagtag  cggagactac  aggcgcctgc  caccacaccc  agctaaattt  gttcttttct
1621 ttctttttt  ttttttggg  gatttttaga  cagacgggg  ttcatatgtt  ggccaggatg
1681 gtctcgatct  cttgacctgc  tgatccgccc  gcctcagctt  cccaaagtlc  tgggattatg
1741 ggctgtagcc  actgcactag  gcctaatttt  tttattttta  gttagatgg  ggtttacca
1801 tgttggccag  gctggtctgg  aacctctgac  ctcaagtgg  ctgccctctc  cagcctccca
1861 aagttctgag  attacaggca  tgagccattg  cgtctgaccc  agactcctta  atgtgactaa
1921 ctccaggctt  tcttggact  acttcttact  tgtctttcca  gctttgtctt  ttcacctctc
1981 caattgagat  aaaataataa  caacctcttg  gatttctcat  caggattaca  tgaaatgaga
2041 tatgtaacat  gcttagcagt  gcctgtccat  agtaaatctc  aataaatgtt  tgtggaatta
2101 taatatcttg  tcatgtttga  gactttgtct  tgcataatca  ggcaccagta  ggtttttata
2161 aaggaaccog  tctgtcacgt  gcagaggaga  aataaacaga  aagtttccca  tctcaggga
2221 gccacctgac  tgacagaggc  acagtgcac  cactctccag  gtctagggga  gaaagcagcc
2281 ttatttctta  gtagctcaga  atctgacttg  agaaacacat  ccacatagaa  aaaaacaagg
2341 aactttttcg  ggtcagggtc  cgggacccac  agtgagggtg  aagatacagg  ggaagggaaga
2401 gggaaataga  gccatcccca  ggggtgaaga  tctcagaaga  gaatttggga  aacaagggtat
2461 gaacaaggac  tgaatagtga  gaagtgtatg  agagacagct  aaagtagatg  gagtgtcaaa
2521 accaaaacct  ctaagggtag  aataggcagc  aatttggcca  agtcctaaca  gggaggccca
2581 taggaggatt  caacctcaag  atgctgtgcc  acattccaag  agggaaacct  aaggctgggc
2641 tgaagagtca  gagatggcta  cagctggcaa  aaagatgggc  agatgctgag  aggagatgat
2701 tgctaaaaatg  ttctgtccag  gacattcaca  gtatctctat  aaccagagtc  tttttgtctg
2761 ttgttgttct  caagaaggaa  acttgaggcc  ggggtgtggtg  gtttatgccc  ataattccag
2821 cgctttgggg  ccaaggcagg  cggatcacct  gaggtcagga  gttcgagacc  agcctggcca
2881 acagtgtgaa  acctcatct  tactaaaaat  acaaaaatta  gctggatgag  gcggtagggtg
2941 cctgtaattgc  cagctactcg  ggagctgag  gcaggagaat  cacttgaacc  tggaggcgcg
3001 aggttgcagg  gaggcggagg  ttgcagtga  ccaagattgc  accactgcac  tccagcctgg

```

FIG. 18

3061 ggcacagaga gtaagactgt ctcaaaaaat aaatgaataa ataaaaagga agaagaagaa  
3121 gaagaacaat tgcaatcctc cctggctcta gaatgtcatt taaaagtcga gtgtcttctt  
3181 ccttccctgt tttgaagcag ccttctctat gacaggcttg cttgccagg ttccctctga  
3241 ccttaaatct ctctcttttg gtgtcttggg cagggcagtt cagaqtgata ggaccaagac  
3301 accctatccg ggctctgggc ggggatgaag tggaaattgcc atgtcgcata tctcctggg  
3361 agaacgctac aggcattggg gtgggggtgt accgcccccc ctctcttagg gtggttcctc  
3421 tctacagaaa tggcaaggac caagatggag accaggcacc tgaatatcgg gcccgacag  
3481 agctgtgaa agatgtctatt ggtgagggaa aggtgactct caggatcccg aatgtazggt  
3541 tctcagatga aggaggtttc acctgcttct tccgagatca ttcttaccaa gaggaggcag  
3601 caatggaatt gaaagtagaa ggtgagtagt gccatataat attaggtatt aactgttggg  
3661 tggccaagaa caattattct ctcaactgag atgagatccc tcaacccaaa catctcagtc  
3721 ctgggaatga tttccataaa aatgtacaca tcaataaaca gaaactcatg cttagggatg  
3781 tctgttgcatt cattattcag agtagcaagg aaattgggat caaatcaat gcctttgagt  
3841 aggtaagtga cagaatgaac aatggtagcc atactgtgaa tattatgcag ggattaaaaa  
3901 gcttattttt gcactaggcc agatggtttg gggggctcct ctaagggtatt attgaggtat  
3961 aagagcaagc tgctgtagga taaaaaaca aaaacaaaac cctagggcat ggtggtttgc  
4021 ctgcagccta ctcaggaggc tgagacggga ggctggcttg agcccagggg tttgcagtta  
4081 cagtgtgcta tgattgcacc actgcactcc aaccgggtg acagagcaaa gaccttcacc  
4141 cccactccct acccgctctc aaaaaaaca aaaacaaaaa caaaaaaacc cttgggcccc  
4201 gcgcctgtgc tcacgcctgt aatcccagca ctgtgggagg ccgaggtggg cagatcacaa  
4261 ggtcaggaga tcgagaccat cctggctaaa acggtgaaac cccgtctcta ctaaaaatc  
4321 aaaaaaazaa aaaaaattta gccaggcatg gtagcaggcg cctgtagtc cagctactcg  
4381 ggaggctgag gcaggagaat ggcgtgaaac cggaagcggg ggttgacag agccaaaatc  
4441 cttccactgc actccagcat ggtgagcgca cacacacaca cacacctgtg cttggtccta  
4501 accctgtatt tgtgagcgca aatgtctaaa tataattata gaaaggagat gtcacctttt  
4561 gtgaataagc aagtaaatca aatgtctaaa tataattata gaaaggagat gtcacctttt  
4621 ggctgtacct ccaactatttc attctgcaga attgcagaat ttcttttttt ttctcttct  
4681 ttcttttctt ttcttttttg acacagagtc tcgctctgta acccaggctg gagtgcattg  
4741 gcgcctcccg cctcctgggt tcaagtgtt ctctgcctc agcctcccg gtagctggga  
4801 ttacagggtc ccaccaccac acccagctaa tttttgtatt tttagtagag acagggtttc  
4861 accagggtgt caagggttgt ctcaactcc tgacctcagg tgatccactc gctcagact  
4921 cccaaagtgc tgggattaca ggcattgagc atggtgccc gctcagaat ttcatattca  
4981 acatgttttg catgatgggt gattttggag aatatttttt gctctatcgc aggatgatta  
5041 agatgtggac aaggtgaagc cgatggaggg ggagctttga aagttacttg ctatttaatt  
5101 gaggaactaa actgctttga gagcctgggg gtcagatcct ctgcttttc cctcccca  
5161 cctgcagtcg aaacatcaga caattgtatc ctattgtatc ttggagggtg gagtgcattc  
5221 tgcagtgctg ggaccagaag atggcattgt atgtggaaca acaaagcact atttctagag  
5281 actgcctgca gggatatgga aatagcttta tgtgtctcag aatgttcttc atacagctgt  
5341 ttttatttgg gaaattctac ttgcccgaaa gtttgatagt gagacctct ccagtttgca  
5401 gatttttctc ctctctgctc aacaacttcc tagctcagta actgcctctc ccaacaaact  
5461 cctcagttt caccacacca zaaaagggaag acaagccggt tgcggtggct cacacctata  
5521 atcccaaaac tttgggaggc cgaggcgggt ggatccacct gaggtcggga gttcgagact  
5581 agcctgacca acatggagaa accctgtctc tactaaaaac acaaaattag cctggcgtgg  
5641 tggcgcattc ctgtaatccc agctgggagg ctgaggcagg agaactcgtt gaaccccgga  
5701 ggcggagggt gcagtgagcc aagatcgttc cattacactc cagtctgggc aagaaaaagt  
5761 gaactccatc tccaaaaaaa aaaaaaaaaa zacaeggaag zcazaaagea aagcagctaa  
5821 agactttgac tcaggggaga aagttctctt ttgggttgct atccacattc caacctcctg  
5881 tttccacactc ttctgtctga tgcttagaa actgttttac aagtaataaa gggacgcttt  
5941 gcttaggctt tggagccagg aagttgagac zaattttaga atcagatgaa gtaattggtat  
6001 tattgcaagt ctcaggtgta actacctctg ctcttctctc gaagagtttc taatttctct  
6061 tgtttactta ttttttctt gtcatttttg ggattttatt actagttgtc tctaactcct

FIG. 18 (cont.)

6121 tcttttaaatt cttcattatg aaacataaaa acaaatgcc a ggcgcggcag ctcacgcctg  
6181 taatcccagc actttgggag gccgaagcgg gcagatcacc cgggtcagga gttcagagacc  
6241 agcctgatca acatggagaa acccgcgtct tactaaaaaa tacaaaatta gctaggcgctg  
6301 gtggcacatg ccagtaatcc cagctacttg agagactgag gcaggagaat cgcttgaacc  
6361 gggaggcaga ggttgcggtg agccaagatc gcgccattgc actccagcct gggcaacaag  
6421 agcaaaactc tgtctcaaaa aaaaaaaacc acatacaaac cagagataat attataatga  
6481 gcctccaagt gcctaccacc ttgtgcagc acttgtcaat ccagggacca cccacctcac  
6541 cggctcccca ctcattacca cctccccc ta ctaattact gaggtaaatc ctaggcagca  
6601 tgatcatttc ttttttttct ttttatttat tttgagacag gatctgtctc tgtcaccag  
6661 gctggagtgt agtggcatat ctctgctcac tgcagcctct gcctcccggg cagaagccat  
6721 cctccacact cagcctacat agtagctggg acccacaggc cacaccacca cacactgcta  
6781 atgttttcta tttttttag agactgggtt ttaccatgtt gatcaggctg gtctcaact  
6841 cctaggctca agcaatctc ccacctcggc ctcccaaagt gctagaatta caggcgcgag  
6901 ccactgcacc cagcgaagaa cactttttaa aaaataaata ggcggggcgc ggtggctcac  
6961 acctgtaate ccagtacttt gggagcccaa ggaggcgaa tcatgaggtc aagagattga  
7021 gaccatccta agtaacatgg tgaacccca tttctactac aaatacaaaa acaaaattag  
7081 cctggcgtgg tggcaggcgc ctgtagtccc agctacttgg gagctgaggc aggagaatgg  
7141 agtgaaccog ggaggcggag cttgcagtga gctgagatca tgccactgca ctccccctg  
7201 gggcaacaga gtgagactcc caaaaaaaa aaaaaaagcc cccctcccc acacacaata  
7261 atataaataa ataaataacc acaatactat tatcacatct taaaaactca acaaaatatt  
7321 cttaatatca tcaaatatcc agtttgtgtt caaattttcc tgattgtttc ataaatatac  
7381 tcttacagtt ggtttctttt agcgagattc aaatgagacc cactgttga cctttgccct  
7441 tagggtttcc caggggtctga attttgttga cgacattccc atgttgcctat gtaatacggg  
7501 cctccatgcc ctgtgttttt ctgtaaactg atagatgtgg aggtcgaatc acatttgtgt  
7561 ttgatttact ttggcaata tagttcatca gtgatactct atacttcttg ttgctttaca  
7621 tccggagggt gataatgtct gcttttctct cttttcta at ttttgtga aggaataatg  
7681 tgggggggtg ggagaaaaaa acccttaagt acatactcgc taaatcacat tgctacaggc  
7741 aacttccatt aagaacttga aagtaaagggt agctgcattt tcccctaggg aacacaatga  
7801 tagacaggag ccttagtcta cagcttgaag gattgtaatt atacctaac aacctcctg  
7861 gaccagttta atgttattag ctgtgatgta tccctacctt tgatgtcatt atccttactt  
7921 agtcccttta aagcagagat caagatgaaa agggcttcag ctgcagcatg gcacatggag  
7981 attagagtgg ggcttttggg tgtgaggag cagacctaga atgggaaata gatgggagcc  
8041 acagaagtga aggtccccc ctctcattgc tcaacctact ccacatctcc aggtctgcac  
8101 atctgttcag ttactgaatc ctgtgtaagc taccttcttt ttcttttttc ttttatttat  
8161 ttatttattt tttttttgag atggagtttt gctcttgtta cccaggctgg atgcaatgg  
8221 tgcaatctcg gctcactgca cctcccaact cccaggttca tgcaattctc ctccctcagc  
8281 cttccaagta gctgggatta caggctgcac caccatgtct ggctaatttt tgaaaaatca  
8341 gtagagagag ggtttcacca tgttggccaa gccggtctcg aactcctgac ctcaagtga  
8401 ccacccacct tggcctccca aaatgctggg attacagggt tgagccacca tgcccgctgt  
8461 aaactacctt cttaaaagct ctagaagagg gcttttaacc ttttgttgtg tgtcatgcac  
8521 cttccgcaag ctgatgaagt tgatagacc atctcagaat tttttttttt tttttgagac  
8581 agtgtctcac tctgtcacc aggatgggtt gcagtggcac gatcatgggt cattgcagcc  
8641 tccacctccc aggtcagat gatcctctg actcagcctc ttgaatagct gagaccacag  
8701 gcttgtgtca ccattgcccag gtaattttta attttttttc gttagggcag ggtctcacat  
8761 tatgttgccc agtctggcct cgagaactcc tgggctcaag caatcttctt gccttgggct  
8821 cccaaagtgg tgggattaca ggggagagcc accacacctt gccaggagga tgttttaaat  
8881 acaccaataa aaacatttat acccaataac agttatccaa atattaaatt aacagaggtt  
8941 aggtgaccc tattaattag tgaatttcc aaatagtaat gaacataagt gatagtttga  
9001 ctttctctg acttttctaa tgtgacgtga aaatatttgt gatttttctt tttctttt  
9061 ttttttgaga tggagtctcg ctcttgttgc ccaggctgga gtgcaatggc aagatctcgg  
9121 ctcacctcaa cctccgcctc ctgggttcaa gcgattctcc tgccctcagcc tcttgagtag

FIG. 18 (cont.)



9181 ctgggattac aggaactgtgc caccacgtcc agctaatttt gtattttttag tagaaacagg  
9241 gtttctccat gttgggtcagg ctgggtcttga actcccaacc tcaggcgatc cgcccgccctc  
9301 ggcctcccaa agtgctggga ttacaggtgt gagccaccgc acctggccaa tatttgtgat  
9361 ttttattgac gacaaagtca aaggttctct tcatattatt gtggtgtatc gcctaccagc  
9421 ataattaaaa taaacactaa atttcagttt aaagtttact gaaaaataat atgtattttt  
9481 tattccctat ttaagctttg aatccctga ctccctatac cattaccact gtcctagttc  
9541 aggttcatgt tgttttttac ttttaattgtt atcacagtct cttaacattt ctcctatgt  
9601 tctccagtc tgtaggtgct aaatctgacg tggtcacttc tcagcttggg atccttcagt  
9661 gcaccaccac agccttgaac tacatatttg aaatacatat ttattttcag taaactttaa  
9721 actgaaattt agtggtttatt ttaattatgc ttgtaggcga tacaccacaa taatatgaag  
9781 agaacctttg actttgtcgt caataaaaag tcccttgagg ggacttcaga tgaagtccc  
9841 ttagctgtct gttaaaaact cccagggctg acccaatata caatcttgac tttaaaccac  
9901 ttgtcattct aaatcactag catctcctgg aaaaaaagc cattttctct tcagggtcaa  
9961 gctcagggac caattctgtg tcaccttctt tgaatcctga tgatattcac ttctttattt  
10021 gacctgattt attgggcccc agacaccatg ctgagtgttg gggattcagc tctggacaat  
10081 gtcaaatgtc agtccctgctt ttcagatcct ttctactggg tgagccctgg agtgctgggt  
10141 ctctctcggg tgcctgctgt gctcctcctg cagatcactc ttggcctcgt cttcctctgc  
10201 ctgcagtaca gactgagagg tacagggcag aggtgtgggt gatcaggatc cttctcttaa  
10261 atgagctggc ttcttgaggc tacaccactt aacatgtatt tgtgagtgc ttctgggttc  
10321 agaagtctct ctactattg agtgataaag aaaaaaata actccatgat gaaagagttt  
10381 tacatcttac ggaatgcttt catatgaata atcggacctg gcatttccct atgagctaac  
10441 tatgcatat agtaacccca ttttacagag gatacaactg aggccaggag tagttcagt  
10501 acttactcaa accgatataa cttataagtg gttagagctga ggccctctga tcataacctg  
10561 cagctccatg caecttgga gagtgtgagc ttcgaagtca gacaggtcta ggctattagg  
10621 agttttgaat aaagatactg aagtgaagc ctctaccaca cagtaggcct tcgaaaattg  
10681 tttcctcttt ctccattcaa cactgaggac tcagggtcag ctgctgatga agctcctctt  
10741 ttttgcttag agctttcatt ctgagccttc tctcctacc aagtgctctc ccaatgccag  
10801 agcaggaaga gtcttcactc ctcccaatgc cccacctccc atttgttact aagaggagag  
10861 gagaaagttag caaggagggt atgggggaatg ttctggggga atgggtgttg gtgcgatcaa  
10921 caacaaaagtc ctttctctca ccttgaattc atcccagatg cctgcttgtt tacttcttcc  
10981 acacaaaaaa aggccttcag cctcatggc tgagcagaaa gaatctgaat gttagagtca  
11041 ggcagcctgg gtttgaattc catctcaggt actgaactct atagcaaaat tcttagattc  
11101 tccaagcttc agttgccttg tctgtcaaat agagaaaaa tccctcctcc taaattgtag  
11161 ggaggattaa agtcatgcaa agtgctcact acaaattccag tcacaaagta gctagctact  
11221 cactaaatgt tcagctcctc cctcctcatt cagatgggga gtggttttag ataaacaaag  
11281 tggcaacgca gtgggctgga gcagctctgt gaactgagaa tccaagaaaa gggcggaaga  
11341 gcagctggga tgtattggat gcttgtgctg gcttggagca ttgctccat tctttattcg  
11401 ctattgtatc tagactatag cttagaaaa agccgcaacc attggcttta aatccagtgc  
11461 tcttctact ctcttgaggt tgtttccagg ctgcagagaa atagcctgca caaggggccc  
11521 aggcgctggg tgtgggaggg tccccaccga gagccagaac atgcaggaa ctaaatgttg  
11581 cctttttcta ttttaggaaa acttcagagc gagataggtg agttccagtc atcgtttctc  
11641 ccaattcttg ccttttggtt ttttggcata acggaaatgg tccattctt ggaccgtctc  
11701 tccctctcaa taccctgttt tccctcagt tccctttct ctacagtggg tgtgtcgtgc  
11761 ctagaacaag ttttaagtaa ttaataaca aagactcagg ataaaaggat cctttttgga  
11821 gtgcccact aaatccattt ccatttggtt ctctttcaga gaatctccc cggacttttg  
11881 gtaagtctcg gcattgtctg gccctcccag gtcaacttgg tatttctact tagttccagt  
11941 caccctgggg aacaaggacc cctggctcct ggttgagtc cttctctct tctctttct  
12001 tcttttaaat aagaagtcatt ttgcatttag gatttggttaa atcataataa aaatactcat  
12061 gtactgtttt tatgtgccag gcactattct aactacttta caaaaacgtt atcttattct  
12121 gtttaactcc ttatgcacat gatctctctt ttcagggaatg ccaaaacaga ggtaaataga  
12181 tctgtttacac gtaaacctga tgtctggttg gggaggtgaa acaaacagaa acaagacaca

FIG. 18 (cont.)

12241 actgtatcac ctgtacttat atttctgctt tacaaactca ggatgtttcc atgagtacag  
12301 aacatgacta atcagagaag acctcataga ggaatagaaa agccaccaag ccccaactagg  
12361 aattgacccc tcaaggacat gggttctagc ctttttggtc actgcagatt gcccaatgcc  
12421 taaagataat ggcaacagaa gagcacccaa atatttggtt gataaatggt gcagacacta  
12481 gaaggtgtca ttagggcaca gatggtacct tctctgagca aacttccttc acagctcctc  
12541 ctcccagaggc tgtagggtgac tctactcttg tcacctggca cacagagttc tatcgtacga  
12601 tttaggaaat tagaccagtg tgtggaccac acacacacac atctttacac acccaaaagag  
12661 gaggaatagt atctttggtt tggaggactt gactatgaaa ggtcttaact cttttttgta  
12721 ccatgaatct ctctggcact ccagtgaagt cttaaaggacc cctttgcaga atgtttttaa  
12781 atatacacat aaaatagaac acataggatt gcaaaaacaa tcattgtact aaaatacagt  
12841 tatcaaccga taatcacatt tgtgatatag taacataaat gtttcttttt ttttttttg  
12901 gaggcagagt ttggtctctg tcaccaggc tggagtgcga tggcgcgac taggctcact  
12961 gaaacctctg cctcccgggt tcaagcgatt ctacgcctcc tgagttagctg ggattacagg  
13021 tgcccgccac cacaccagc taatttttgt atttttagta gagactagggt ttcaccagggt  
13081 tggccaggct ggctcgaac tctgacctc aggtgatcca cctgccttg cctcccaaag  
13141 tgtctgggatt acgggcata gcccctgctc ccggccataa atatttcttt agccaaagta  
13201 atacattaag taatgtagca gcaagtctaa taacctgtaa tttctttctt tcttctttc  
13261 tttctttttt tttgagatga agtttttttg agatggagtg caatggcaca atctcggctc  
13321 actgcaacct ccacctcctg ggttcaagcg attctcctgc ctacgcctcc caagtgtctg  
13381 gaactacagg cgcattgccac catgcccagc taatttttgt atttttagta gagacgggggt  
13441 ttcaccatgt tggccaggct ggtcttgaac cctgacctc aggtgatctg cctgccttg  
13501 ccttccaaag tgcctgggatt acaggcatga gccaccaggc ccagccaat aacctttaat  
13561 ttcaacatac taataaacat aaacgtatt tcaagatttc tgcaataact ctaatgggaa  
13621 tgaazacatc tgtggcttcc attggttaatt aagtcacagg tactgctcat atgtgtgta  
13681 gttgtaaaaat gtttttggtt gttttgtttt ttccaagact tgggggaatg ggtgttggtg  
13741 ggatcaacaa gegtcttctt ctgtggccca ggctggagtg caggggcagg atcttggtc  
13801 actgcaacct ccgcctcca ggttcaagcg attctcctgc ctacgcctcc tgagtactg  
13861 gcattacagg catgtgccac cagcccagc taatttttac atttttagta gagatgggggt  
13921 ttcacatgt tggcctggct ggtcttgaac tcttggtcctc atgatccacc cgtctcggac  
13981 tcccagagtg ttgggattac aggcattgagc caccacacct ggcagttgtt acatttttaa  
14041 tgaagaaaaa tgttaaatcc agttattgaa aataaggagg cagtactttt ctcatccaa  
14101 ttcattgagt ttctgaattt tgtcccccga gtcttttgtt gttctaggac cccaggttaa  
14161 ggaacccaaa aagacagggt ggtggggcat gagggggaac acatgttaat ccctgttgt  
14221 tctggtgaac aattcagatc cccactttct gaggggtgcc tgccttgatc atctgctaca actggctaca  
14281 tgtaattgtg ccggttcttg gaccttgggt tgccttgatc atctgctaca actggctaca  
14341 tcgaagacta gcagggtgcag tggctgggca gcaggcaaga ccaccaataa gtgggggacc  
14401 aagtcagctc tgaatgggaa gccaaaagag aatagaacca ggactcaaga ttaggggagc  
14461 tgggattttc ttattcctct gtccccatgc ccaaccccag gctcttctga gaaactgtga  
14521 agagaaccac ttactggatc tgtgggatcc cccagtggaag agggcagtggt gggctactcc  
14581 aaatgtccat agggaggatg tggggaagggt gctattcctc ttccactaat cacatatttg  
14641 tttctttttg ttttcagggc aattccttga agagctacgt aagttctctt ctctctgtta  
14701 taagcagaga ataaaaagcc aggaaggga gacagaagca acaagggaag gaggggggt  
14761 attgagggat cacattccca gaggaaggga ggagctggag agcctgggtg gaggggaagc  
14821 tctcctggg aggttagagg caaagaagcc agctgttaga gacacattta cagggtggcag  
14881 agaagctgga ggcactccta tctgccacct gatccattcc tcttcaactg cccctaagca  
14941 ggaatccaac cctagctggt ctcatgtccc attccacagc aactgcccag tgcctcacct  
15001 ctgagatcaa ccatgtaggc aggaatggag acaagatgac ccaagggtc .tttcttctc  
15061 ctagtccaat ggttttatga tacaactac tgacatacgt ttttcaagct attttctcct  
15121 tcttctagga aatcccttct gagtgatgtc acatcttggc aggggtggag gagagcctgg  
15181 ttgcccaggg atttgtcctt ggggacatct catccatcaa gttgcacact cactggcatc  
15241 tttgctatgg ggacattcca atttgcactt tcaggaaacac tctgaattcc aagtagaatt

FIG. 18 (cont.)

15301 gatttccctt cttctgtcat ctaccttttc tcttcatttt cccattttta ttacccttct  
15361 ttccattttct ctctccagtc ttccacctgg aagccctctc tggctaagga caggcagggtg  
15421 cccctctctc catcagagga cacctgtact ggagagcaac acaggatggg ctctgccatg  
15481 aactggaggc caggaatctc ctactgaaa attacagtat ggtaactttg caaatgggtg  
15541 ttgtttcttc caagactcca gccctgattg cgcaaaactg aaaggcatgt gaagggaagg  
15601 aagaggaaga gtgcaaaaca ttgaagagag agctgagtga gctgaagagt gaggatatga  
15661 gtagcccca cccaaacctg gagatgggga gaaacctaca gaatactagc cagagctcct  
15721 ccttgtcttg gcagcctact agggacctgg ggaagcaaaa acgaaagctg ggcaacatgc  
15781 ctgctttaga atgttttctt tctacttaca catettccac aggtctcaga atctttcctt  
15841 cctctcatcc ttttctccta tctacatata tatcagagta tccactgttt attcaacaac  
15901 tactacttga tggtcagaca caaacaaca agctagggtgc taattaataa agatacaggt  
15961 tttggccggg tgcgggtggt cagcctgtga atcccagcac tttgggaggc cgaggcgggc  
16021 gaatcacgag gtcaggagtt caagaccagc ctggccaaca tgggtgaaacc ccatctctac  
16081 taaaaataca aacaattaac tgagcatagt ggtgggcacc tataatacca gctactccgg  
16141 aggcctgaggc aggagaatcg cttgaaccca ggaggcagag gttgcagtga gctgagatg  
16201 cgccactgca ctctagccgg agtgacagag taagactctg tctcaaaaat aaataataa  
16261 ataaataaat aaataaatpa ataaataaaa aataataata caagttttca taagcacact  
16321 tctaaccctt tgtcttttat gtatttctt ccttatccac gcacctgtct cctctactc  
16381 cagcctcatt accccagagg tcagtcttca ggaaaactaa acacaaagaa agagctcagt  
16441 cagaaaggcc atttatitatt gtttcaagat gctcactgcc tctttgttt tgtctcttt  
16501 gcaggccttc tctcttaggc ctcttctctt gggggtatgg atcctggggg gagattgac  
16561 acctccatgc ttccattcct ccccagccat agtggggaca tcatgagaga agccaagcca  
16621 ctggcccagg atcaccggc atttatgtgt gctgctcttg cacaggctct tgcctttata  
16681 gcccctccag tgatccataa ggccctctt ctcccaaaag gagaggctac agatagggca  
16741 aaggtagctc ttctgtcttc agtgggtctg ctgggtgtctg accagcctgg aaaatgagct  
16801 gaaagacttg ctgcaatgga agcagtagtt gggcggtct gtgagggtgc ccttctggtg  
16861 tctggagaga taggatttct tgctaaaagt caaagaacaa tgggggcaac agaagacatt  
16921 gagtcttgag ggcttctact gatgagagtt ggatctggca tctgacaga gggttccagt  
16981 gatgggtgcc tgggtcctgg tcacagggtgc ttggttctta agtacagatg cctggttctg  
17041 ggccatagga cctcagttc taaatatggg ttcttgggac ctggccactg gtgcatggtt  
17101 cacatccaaa agcccctgga tggacctctg gcttctggcg atgggtgtct ggaattcagc  
17161 ctgggtgcct ggaatcttca aagtaacact ctggtttcca tccactggct cctggttttg  
17221 gtgtatcttc tgggtggcgtt tgagctcaga ctgggtcccgg aagctcttcc cacacacaga  
17281 gcatgaatgg ggccggtaac ccagatggac gcggcggtga cgacttagtc cagaagcatc  
17341 acagttagtc ttgtcacaga gcgtgcaaca gaaggccctc tcccaagat gcattgcgtc  
17401 gtgatactg agggacttgg ggctccgaaa caacttccca cactgactgc agctgttagt  
17461 cagcttggga ttgtgaacaa actggtggct atagaggtag gagcgctgc tgaacattt  
17521 ggcacagggtg tagcaaaa

FIG. 18 (cont.)

```
1  tttgtatgtc attgcaggat tcatgctttc cagtgtgtca tctatggaac tgcctctttc
61  ttcttccttt atggggccct cctgctggct gagggcttct acaccaccgg cgctgtcagg
121 cagatctttg gcgactacaa gaccaccatc tgcggcaagg gcctgagcgc aacggtaaca
181 gggggccaga aggggagggg ttacagaggc caacatcaag ctcatctttt ggagcgggtg
241 tgtcattggt tgggaaaatg gctaggacat cccgacaagg tgatcatcct caggattttg
301 tggcaataac aaggggtggg gggacaa
```

FIG. 19

1 ctgtatcagt gctcctcgtc gcctcactgt acttcacgga agagacttgg ttgactggcc  
61 acttgaggcg gaatcaggag acattcccaa ctgagagaga ctgagcccta gctcgccac  
121 ttgctggaca agatgatatt ccttaccacc ctgctctgtg tttggataat gatttcagct  
181 tctcgagggg ggcactgggg tgcttgatg ccctcgccca tctcagcctt cgagggcacg  
241 tgtgtctcca tccctgccc tttcgacttc ccgatgagc tcagaccggc tgtggtacat  
301 ggcgtctggt atttcaacag tccctacccc aagaactacc cgccagtggg ctccaagtcc  
361 cgcacacaag tggccacga gacttccag ggccgtagcc gcctgttggg agacctgggc  
421 ctacgaaact gcacctgct tctcagcacg ctgagccctg agctgggagg gaaatactat  
481 ttccgaggtg acctgggccc ctacaaccag tacaccttct cggagcacag cgtcctggac  
541 atcatcaaca cccccaacat cgtggtgccc ccagaagtgg tggcaggaac ggaagtagag  
601 gtcagctgca tgggtccgga caactgccc gagctgcgc ctgagctgag ctggctgggc  
661 cactgagggg taggggagcc cactgttctg ggtcggtgc gggaggatga aggcacctgg  
721 gtgcaggtgt cactgctaca cttcgtgctt actagagagg ccaacggcca ccgtctgggc  
781 tgtcaggtgt ccttcccaa caccacctg cagttcgagg gttacgccag tctggacgtc  
841 aagtaccccc cgggtattgt ggagatgaat tctctgtgg aggccattga gggctccac  
901 gtcagcctgc tctgtggggc tgacagcaac ccgccaccgc tgctgacttg gatgcgggat  
961 gggatggtgt tgagggaggc agttgctgag agcctgtacc tggatctgga ggaggtgacc  
1021 ccagcagagg acggcatcta tgcttgcttg gcagagaatg cctatggcca ggacaaccgc  
1081 acggtggagc tgagcgtcat gtatgcacct tggagccca cagtgaatgg gacgggtggtg  
1141 gcggtagagg gggagacagt ctccatcctg tgttccacac agagcaacct ggacctatt  
1201 ctcaccatct tcaaggagaa gcagatcctg gccacggtea tctatgagag tcagctgcag  
1261 ctggaaactcc ctgcagtgc gcccgagcac gatggggagt actggtgtgt agctgagaac  
1321 cagtatggcc agagagccac cgccttcaac ctgtctgtgg agtttgctcc cataatcctt  
1381 ctggaatcgc actgtgcagc ggccagagac accgtgcagt gcctgtgtgt ggtaaaatcc  
1441 aaccgggaa cctccgtggc ctltgagctg ccttcccgca acgtgactgt gaaccagaca  
1501 gagagggagt ttgtgtactc agagcgcagc ggccctcctg tcaccagcat cctcacgctc  
1561 cgggggtcagg cccaagcccc accccgcgtc atttgtacct ccaggaacct ctacggcacc  
1621 cagagcctcg agctgccttt ccaggagaca caccgactga tgtgggcca2 aatcgccctt  
1681 gtgggtgctg tggtcgcctt tgccatcctg attgccattg tctgctacat caccagaca  
1741 agaagaaaaa agaacgtcac agagagcccc agcttctcag cgggagacaa cctcatgtc  
1801 ctgtacagcc ccgaattccg aatctctgga gcacctgata agtatgagag tgagaagcgc  
1861 ctgggggtccg agaggaggct gctgggcctt aggggggaac cccagaact ggacctcagt  
1921 tattccactc cagacctggg gaaacgaccc accaaggaca gctacacctt gacagaggag  
1981 ctggctgagt acgcagaaat ccgagtcagg tga

FIG. 10

1 masqkrpsqr hgs kylatas tmdharhgfl prhrdtgild sigrffggdr gapkrqsgkd  
61 shhpartahy gslpqkshgr tqdenpvvhf fknivtprtp ppsqgkgrgl slsrfswgae  
121 gqrpqfgygg rasdyksahk gfkqvdaqgt lskifklggr dsrsgspmar r

FIG. 24

1 mglleccarc lvgapfaslv atglcffgva lfcgcgheal tgtkliety fsknyqdyey  
61 linvihafqy viygtasfff lygalllaeg fyttgavrqi fgdyktticg kglisatvtgg  
121 qkgrgsrgqh qahslervch clgkwlghpd kityaltvw llvfacsavp vyiylfntwtt  
181 cqsiafpskt sasigslcad armygvlpwn afpgkvcgsn llsicktaef qmtfhlfaaa  
241 fvgaaatlvs lltfmiaaty nfavklmgrp gtkf

FIG. 22

1 maslsrpslp scicsfllll llqvsssyag qfrvigprhp iralvgdeve lpcrispgkn  
61 atgmevgwyr ppfsrvvhly rngkdqgdq apeyrgrtel lkdaigegkv tlrirnvrf  
121 deggftcfr dhsyqeeaaam elkvedpfyw vspgvlvlla vlpvlllqit lglvflclqy  
181 rlrqklraei enlhrtfdph flrvpcwkit lfvivpvlgp lvaliicynw lhrrlagqfl  
241 eelrnpf

FIG. 23